

**Nissl Staining Kit, Toluidine Blue Method****Catalog No.: RA20098****Basic Information**

<b>Product name</b>	Nissl Staining Kit, Toluidine Blue Method
<b>Sizes</b>	100 mL
<b>Storage</b>	RT, keep away from light
<b>Shipping</b>	RT
<b>Validity</b>	12 months

**Product Introduction**

The neuronal cell body includes a large nucleus with a wrinkled nuclear membrane, sparse chromatin, and a prominent nucleolus. In the cytoplasm of the cell body, there are Nissl granules, which represent the rough endoplasmic reticulum and exhibit a characteristic basophilic punctate pattern in many neurons. Nissl granules can be demonstrated by various stains such as neutral red, methylene blue, toluidine blue, and methyl violet. Variations in staining, pH, and differentiation time allow some stains to highlight only Nissl substance, while others also show neuronal nuclei and neuroglia. Nissl bodies (or Nissl granules) are triangular or oval granular structures distributed in the cytoplasm of neurons. They can be stained purple-blue by basic dyes such as thionin, methylene blue, toluidine blue, and cresyl violet. All neurons contain Nissl bodies, but their shape, quantity, and distribution vary. Nissl bodies are also present in dendrites but not in axons or the axon hillock. Their appearance changes with the physiological state of the neuron. Nissl bodies are important sites for protein synthesis; when neurons are stimulated, Nissl bodies in the cell body decrease significantly.

EnkiLife Nissl Staining Solution (Toluidine Blue Method) uses toluidine blue as the core dye. It is suitable for Nissl body staining in paraffin-embedded tissue sections. The presence or disappearance of Nissl bodies is an important indicator of whether neurons are damaged. In cases such as encephalitis, cerebral ischemia, or axonal reaction, Nissl bodies may dissolve or disappear.

**Materials Required (Not Supplied)**

1. Graded ethanol, distilled water.
2. Incubator, alcohol lamp, microscope.

**Nissl Staining Kit, Toluidine Blue Method****Catalog No.: RA20098****Experimental procedure**

1. Fix fresh tissue in ethanol, Carnoy's fixative, or 10% neutral formalin, then proceed with routine dehydration and paraffin embedding.
2. Cut sections at 6–8 µm, deparaffinize with xylene or substitute to water, rinse with distilled water.
3. Immerse sections in Nissl Staining Solution (Toluidine Blue Method), and incubate at 50–60 °C for 25–50 min.
4. Rinse briefly with distilled water.
5. Rinse with 70% ethanol.
6. Rapidly differentiate with 95% ethanol (differentiation is difficult to control; if failed, repeat steps 3–6).
7. Rapidly dehydrate with absolute ethanol, clear with xylene or substitute, and mount with neutral balsam.

**Staining Results**

Component	Color
Nissl bodies	Purple-blue
Nuclei	Brownish red

**Notes**

1. Nissl bodies dissolve easily after tissue removal, so immediate fixation is required; otherwise, staining will be poor.
2. Fixation is critical. Use ethanol, Carnoy's fixative, or neutral formalin.
3. This staining solution works well for paraffin-embedded tissue sections.
4. Rapid differentiation with 95% ethanol is difficult to control; if failed, repeat steps 3–6.
5. Protect stained specimens from light to prevent fading.
6. For your safety and health, wear a lab coat and disposable gloves.
7. Use the reagent promptly after opening to ensure optimal performance.

**This product is for research use only!**