Product Name: LC3A (8F5) Mouse Monoclonal Antibody Enkilife Catalog #: AMM00891

Summary

Production Name LC3A (8F5) Mouse Monoclonal Antibody

Description Primary antibody

Host Mouse
Application WB,IHC-P

Reactivity Human, Rat, Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG1

Clonality Monoclonal Antibody

Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Purification Affinity Purified

Immunogen

Gene Name MAP1LC3A

Microtubule-associated proteins 1A/1B light chain 3A; Autophagy-related protein LC3

A; Autophagy-related ubiquitin-like modifier LC3 A; MAP1 light chain 3-like protein 1;

Alternative Names

MAP1A/MAP1B light chain 3 A; MAP1A/MAP1B LC3 A; Microtubule-associated protein

1 light chain 3 alpha

 Gene ID
 84557

 SwissProt ID
 Q9H492

Application

Dilution Ratio WB: 1/500-1/1000 IHC: 1/50-1/100

Molecular Weight Calculated MW: 14 kDa; Observed MW: 14,16 kDa




Background

Autophagy marker Light Chain 3 (LC3) was originally identified as a subunit of microtubule-associated proteins 1A and 1B (termed MAP1LC3), and subsequently found to contain similarity to the yeast protein Apg8/Aut7/Cvt5 critical for autophagy. Three human LC3 isoforms (LC3A, LC3B, and LC3C) undergo post-translational modifications during autophagy. Cleavage of LC3 at the carboxy terminus immediately following synthesis yields the cytosolic LC3-I form.


Research Area

Signal Transduction

Image Data




Western blot analysis of LC3A (8F5)in Hela lysates, 3T3 lysates, rat Brain lysates using LC3A antibody.



Immunohistochemistry analysis of paraffin-embedded Human Heptacarcinoma Tissue using LC3A antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Immunohistochemistry analysis of paraffin-embedded Human Prostate Carcinoma Tissue using LC3A antibody. Highpressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

Note

For research use only.