

**Product Name: Hsc70 Mouse Monoclonal Antibody**  
**Catalog #: AMM03612**



## Summary

<b>Production Name</b>	Hsc70 Mouse Monoclonal Antibody
<b>Description</b>	Primary antibody
<b>Host</b>	Mouse
<b>Application</b>	WB
<b>Reactivity</b>	Human

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
<b>Purification</b>	Affinity Purified

## Immunogen

<b>Gene Name</b>	HSPA8
<b>Alternative Names</b>	Heat shock cognate 71 kDa protein (Heat shock 70 kDa protein 8)
<b>Gene ID</b>	3312
<b>SwissProt ID</b>	P11142

## Application

<b>Dilution Ratio</b>	WB: 1/500-1/1000
<b>Molecular Weight</b>	Calculated MW: 71 kDa; Observed MW: 71 kDa

## Background

Acts as a repressor of transcriptional activation. Inhibits the transcriptional coactivator activity of CITED1 on Smad-

**Product Name: Hsc70 Mouse Monoclonal Antibody**  
**Catalog #: AMM03612**




mediated transcription. Chaperone. Component of the PRP19-CDC5L complex that forms an integral part of the spliceosome and is required for activating pre-mRNA splicing.

## Research Area

Signal Transduction

## Image Data



Western blot analysis of Hsc70 in HeLa, 293 lysates using Hsc70 antibody.

## Note

For research use only.