

Summary

Production Name	Cytokeratin 18 Rabbit Monoclonal Antibody	
Description	Recombinant Rabbit Monoclonal antibody	
Host	Rabbit	
Application	WB,IHC-F,IHC-P,ICC/IF	
Reactivity	Human	

Performance

Conjugation	Unconjugated		
Modification	Unmodified		
lsotype	IgG		
Clonality	Monoclonal Antibody		
Form	Liquid		
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw		
Storage	cycles.		
Buffer	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05%		
builer	BSA		
Purification	Affinity Purified		

Immunogen

Gene Name	KRT18		
	Cell proliferation inducing gene 46 protein; Cell proliferation inducing protein 46; Cell		
Alternative Names	proliferation-inducing gene 46 protein; CK 18; CK-18; CK18; CYK 18; CYK18; Cytokeratin		
Alternative Names	18; Cytokeratin endo B; Cytokeratin-18; K 18; K18; K1C18_HUMAN; Keratin 18; Keratin		
	D; keratin; type l cytoskeletal 18; Keratin-18; KRT18.		
Gene ID	3875		
SwissProt ID	P05783		

Application

Dilution Ratio	WB: 1/500-1/1000) IHC: 1/50-1/100 IF: 1/50-1/200	



Molecular Weight

Calculated MW: 48 kDa; Observed MW: 48 kDa


Background

K18 a type I cytoskeletal keratin. The keratins are intermediate filament proteins responsible for the structural integrity of epithelial cells and are subdivided into cytokeratins and hair keratins. Keratin 18 and its filament partner keratin 8 are perhaps the most commonly found members of the intermediate filament gene family.


Research Area

Tags & Cell Markers

Image Data




Western blot analysis of Cytokeratin 18 in Hela lysates using Cytokeratin 18 antibody.



Immunohistochemistry analysis of paraffin-embedded Human Cholangiocarcinoma using Cytokeratin 18 antibody.Highpressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.





Immunocytochemistry analysis of Cytokeratin 18 (green) in hela using Cytokeratin 18 antibody, and DAPI(blue)

Note

For research use only.