

## Summary

<b>Production Name</b>	HMGB1 Rabbit Monoclonal Antibody
<b>Description</b>	Recombinant Rabbit Monoclonal antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC-F,IHC-P,FC
<b>Reactivity</b>	Human, Mouse, Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal Antibody
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
<b>Purification</b>	Affinity Purified

## Immunogen

<b>Gene Name</b>	HMGB1
<b>Alternative Names</b>	HMGB1; HMG1; High mobility group protein B1; High mobility group protein 1; HMG-1
<b>Gene ID</b>	3146
<b>SwissProt ID</b>	P09429

## Application

<b>Dilution Ratio</b>	WB: 1/500-1/1000 IHC: 1/50-1/100 FC: 1/50-1/100
<b>Molecular Weight</b>	Calculated MW:25 kDa;Observed MW: 29 kDa

## Background

**Product Name: HMGB1 Rabbit Monoclonal Antibody**  
**Catalog #: AMRe04024**




High mobility group (HMG) proteins 1 and 2 are ubiquitous non-histone components of chromatin. Evidence suggests that the binding of HMG proteins to DNA induces alterations in the DNA architecture including DNA bending and unwinding of the helix. HMG proteins synergize with Oct-2, members of the NFκB family, ATF-2 and c-Jun to activate transcription.

## Research Area

Epigenetics and Nuclear Signaling

## Image Data



Western blot analysis of HMGB1 in HeLa lysates using HMGB1 antibody.

## Note

For research use only.