

Summary

Phospho-Tau (S404) (13R18) Rabbit Monoclonal Antibody
Rabbit Monoclonal Antibody
Rabbit
WB,ELISA
Human, Mouse, Rat

Performance

Conjugation	Unconjugated
Modification	Phospho Antibody
lsotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	МАРТ
Alternative Names	MAPT; Microtubule-associated protein tau; MTBT1; Neurofibrillary tangle protein;
Alternative Names	Paired helical filament-tau; PHF-tau
Gene ID	4137.0
SwissProt ID	P10636.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	79kDa




Background

Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity (PubMed:21985311). The C-terminus binds axonal

microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both (PubMed:21985311" target="_blank">21985311" target="_blank">21985311" target="_blank">21985311, PubMed:32961270). Axonal polarity is predetermined by TAU/MAPT localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization.

Research Area

Image Data



Western blot analysis of Phospho-Tau (S404) expression in mouse brain cell lysate.

Note

For research use only.