

**Product Name: Aurora A (8C5) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe07374**



## Summary

<b>Production Name</b>	Aurora A (8C5) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	AURKA
<b>Alternative Names</b>	AIK, ARK1, AYK1, Aurora-A, Aurora-related kinase 1, BTAK, IAK1, Ipl1- and aurora-related kinase 1, STK15, STK6, Serine/threonine kinase 15
<b>Gene ID</b>	6790.0
<b>SwissProt ID</b>	O14965.

## Application

<b>Dilution Ratio</b>	WB 1:500-1:2000
<b>Molecular Weight</b>	46kDa

**Product Name: Aurora A (8C5) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe07374**



---

## Background

Mitotic serine/threonine kinases that contributes to the regulation of cell cycle progression. Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis. Mitotic serine/threonine kinase that contributes to the regulation of cell cycle progression (PubMed: [26246606](http://www.uniprot.org/citations/26246606), PubMed: [12390251](http://www.uniprot.org/citations/12390251), PubMed: [18615013](http://www.uniprot.org/citations/18615013), PubMed: [11039908](http://www.uniprot.org/citations/11039908), PubMed: [17125279](http://www.uniprot.org/citations/17125279), PubMed: [17360485](http://www.uniprot.org/citations/17360485)). Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis (PubMed: [26246606](http://www.uniprot.org/citations/26246606), PubMed: [14523000](http://www.uniprot.org/citations/14523000), PubMed: [14523000](http://www.uniprot.org/citations/14523000)). Required for normal spindle positioning during mitosis and for the localization of NUMA1 and DCTN1 to the cell cortex during metaphase (PubMed: [27335426](http://www.uniprot.org/citations/27335426), PubMed: [27335426](http://www.uniprot.org/citations/27335426)). Required for initial activation of CDK1 at centrosomes (PubMed: [13678582](http://www.uniprot.org/citations/13678582), PubMed: [15128871](http://www.uniprot.org/citations/15128871), PubMed: [15128871](http://www.uniprot.org/citations/15128871)). Phosphorylates numerous target proteins, including ARHGEF2, BORA, BRCA1, CDC25B, DLGP5, HDAC6, KIF2A, LATS2, NDEL1, PARD3, PPP1R2, PLK1, RASSF1, TACC3, p53/TP53 and TPX2 (PubMed: [18056443](http://www.uniprot.org/citations/18056443), PubMed: [15128871](http://www.uniprot.org/citations/15128871), PubMed: [14702041](http://www.uniprot.org/citations/14702041), PubMed: [14702041](http://www.uniprot.org/citations/14702041), PubMed: [11551964](http://www.uniprot.org/citations/11551964), PubMed: [15147269](http://www.uniprot.org/citations/15147269), PubMed: [15147269](http://www.uniprot.org/citations/15147269), PubMed: [15987997](http://www.uniprot.org/citations/15987997), PubMed: [15987997](http://www.uniprot.org/citations/15987997), PubMed: [17604723](http://www.uniprot.org/citations/17604723), PubMed: [17604723](http://www.uniprot.org/citations/17604723), PubMed: [18615013](http://www.uniprot.org/citations/18615013), PubMed: [18615013](http://www.uniprot.org/citations/18615013)). Regulates KIF2A tubulin depolymerase activity (PubMed: [19351716](http://www.uniprot.org/citations/19351716), PubMed: [19351716](http://www.uniprot.org/citations/19351716)). Important for microtubule formation and/or stabilization (PubMed: [18056443](http://www.uniprot.org/citations/18056443), PubMed: [18056443](http://www.uniprot.org/citations/18056443)). Required for normal axon formation (PubMed: [19812038](http://www.uniprot.org/citations/19812038), PubMed: [19812038](http://www.uniprot.org/citations/19812038)). Plays a role in microtubule remodeling during neurite extension (PubMed: [19812038](http://www.uniprot.org/citations/19812038), PubMed: [19812038](http://www.uniprot.org/citations/19812038)).


**Product Name: Aurora A (8C5) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe07374**

---

<http://www.uniprot.org/citations/19668197> target="\_blank">19668197</a>). Also acts as a key regulatory component of the p53/TP53 pathway, and particularly the checkpoint- response pathways critical for oncogenic transformation of cells, by phosphorylating and destabilizing p53/TP53 (PubMed:<a href="http://www.uniprot.org/citations/14702041" target="\_blank">14702041</a>). Phosphorylates its own inhibitors, the protein phosphatase type 1 (PP1) isoforms, to inhibit their activity (PubMed:<a href="http://www.uniprot.org/citations/11551964" target="\_blank">11551964</a>). Necessary for proper cilia disassembly prior to mitosis (PubMed:<a href="http://www.uniprot.org/citations/17604723" target="\_blank">17604723</a>, PubMed:<a href="http://www.uniprot.org/citations/20643351" target="\_blank">20643351</a>). Regulates protein levels of the anti-apoptosis protein BIRC5 by suppressing the expression of the SCF(FBXL7) E3 ubiquitin-protein ligase substrate adapter FBXL7 through the phosphorylation of the transcription factor FOXP1 (PubMed:<a href="http://www.uniprot.org/citations/28218735" target="\_blank">28218735</a>).

## Research Area

## Image Data



Western blot analysis of Aurora A expression in HepG2 cell lysate.

## Note

For research use only.