

Product Name: Bcl10 (9M3) Rabbit Monoclonal Antibody
Catalog #: AMRe07497

Summary

Production Name	Bcl10 (9M3) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	BCL10 {ECO:0000303 PubMed:9989495, ECO:0000312 HGNC:HGNC:989}
Alternative Names	BCL10; CARMEN; CIPER; CLAP; c-E10; mE10;
Gene ID	8915.0
SwissProt ID	O95999.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	26kDa


Background

Promotes apoptosis, pro-caspase-9 maturation and activation of NF-kappa-B via NIK and IKK. May be an adapter protein between upstream TNFR1-TRADD-RIP complex and the downstream NIK-IKK-IKAP complex. Is a substrate for MALT1. Plays a key role in both adaptive and innate immune signaling by bridging CARD domain-containing proteins to immune activation (PubMed: [10187770](http://www.uniprot.org/citations/10187770), PubMed: [10364242](http://www.uniprot.org/citations/10364242), PubMed: [10400625](http://www.uniprot.org/citations/10400625), PubMed: [25365219](http://www.uniprot.org/citations/25365219), PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Acts by channeling adaptive and innate immune signaling downstream of CARD domain-containing proteins CARD9, CARD11 and CARD14 to activate NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Recruited by activated CARD domain-containing proteins: homooligomerized CARD domain-containing proteins form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10, subsequent recruitment of MALT1 and formation of a CBM complex (PubMed: [24074955](http://www.uniprot.org/citations/24074955)). This leads to activation of NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed: [18287044](http://www.uniprot.org/citations/18287044), PubMed: [27777308](http://www.uniprot.org/citations/27777308), PubMed: [24074955](http://www.uniprot.org/citations/24074955), PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Activated by CARD9 downstream of C-type lectin receptors; CARD9-mediated signals are essential for antifungal immunity (PubMed: [26488816](http://www.uniprot.org/citations/26488816)). Activated by CARD11 downstream of T-cell receptor (TCR) and B-cell receptor (BCR) (PubMed: [18264101](http://www.uniprot.org/citations/18264101), PubMed: [18287044](http://www.uniprot.org/citations/18287044), PubMed: [27777308](http://www.uniprot.org/citations/27777308), PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Promotes apoptosis, pro-caspase-9 maturation and activation of NF-kappa-B via NIK and IKK (PubMed: [10187815](http://www.uniprot.org/citations/10187815)).

Research Area

Image Data

Product Name: Bcl10 (9M3) Rabbit Monoclonal Antibody
Catalog #: AMRe07497



Western blot analysis of Bcl10 expression in HeLa cell lysate.

Note

For research use only.