Summary

Production Name C9 (12U11) Rabbit Monoclonal Antibody

Description Rabbit Monoclonal Antibody

Host Rabbit
Application WB,ELISA
Reactivity Human

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type
Buffer	preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term.
	Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name C9

Complement component C9; Comple Complement component C9bment component Alternative Names

C9a;

Gene ID 735.0 **SwissProt ID** P02748.

Application

Dilution Ratio WB 1:500-1:2000


Molecular Weight 63kDa

Background

C9 is synthesised in the liver and monocytes, and is a plasma protein consisting of a single polypeptide chain of molecular weight 71kDa. Normal plasma concentration is 60mg/L. C9 forms part of the membrane attack complex (MAC) the cytolytic terminal complex of the complement pathways. C9 binds to the membrane associated C5b-8, binding of C9 to C5b-8 leads to the circular polymerisation of 12-18 C9 molecules. This is the basis of the hydrophilic, protein-walled, trans-membrane channel formed by the MAC, which leads to cell lysis and destruction. Constituent of the membrane attack complex (MAC) that plays a key role in the innate and adaptive immune response by forming pores in the plasma membrane of target cells (PubMed:9634479, PubMed:9212048, PubMed:26841934). C9 is the pore-forming subunit of the MAC (PubMed:26841934, PubMed:26841934).

Research Area

Image Data



Western blot analysis of C9 expression in Human fetal liver lysate.

Note

For research use only.