

**Product Name: CD14 (16H16) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe08209**

---

## Summary

<b>Production Name</b>	CD14 (16H16) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	CD14
<b>Alternative Names</b>	CD14; Monocyte differentiation antigen CD14; Myeloid cell specific leucine rich glycoprotein;
<b>Gene ID</b>	929.0
<b>SwissProt ID</b>	P08571.

## Application

<b>Dilution Ratio</b>	WB 1:500-1:2000
<b>Molecular Weight</b>	40kDa

**Product Name: CD14 (16H16) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe08209**




## Background

CD14 antigen is a GPI-linked glycoprotein with a molecular weight of 55kD. The CD14 antigen is expressed on cells of the myelomonocytic lineage including monocytes, macrophages and Langerhans cells. Low expression is observed on neutrophils and on human B cells. CD14 antigen is a receptor for bacterial lipopolysaccharide (LPS, endotoxin) and the lipopolysaccharide binding protein (LBP). LBP and CD14 antigen serves two physiological roles. Coreceptor for bacterial lipopolysaccharide (PubMed: [1698311](http://www.uniprot.org/citations/1698311)), PubMed: [23264655](http://www.uniprot.org/citations/23264655)). In concert with LBP, binds to monomeric lipopolysaccharide and delivers it to the LY96/TLR4 complex, thereby mediating the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed: [20133493](http://www.uniprot.org/citations/20133493)), PubMed: [23264655](http://www.uniprot.org/citations/23264655) target="\_blank">23264655</a>, PubMed: [22265692](http://www.uniprot.org/citations/22265692) target="\_blank">22265692</a>). Acts via MyD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed: [8612135](http://www.uniprot.org/citations/8612135) target="\_blank">8612135</a>). Acts as a coreceptor for TLR2:TLR6 heterodimer in response to diacylated lipopeptides and for TLR2:TLR1 heterodimer in response to triacylated lipopeptides, these clusters trigger signaling from the cell surface and subsequently are targeted to the Golgi in a lipid-raft dependent pathway (PubMed: [16880211](http://www.uniprot.org/citations/16880211) target="\_blank">16880211</a>). Binds electronegative LDL (LDL(-)) and mediates the cytokine release induced by LDL(-) (PubMed: [23880187](http://www.uniprot.org/citations/23880187) target="\_blank">23880187</a>).

## Research Area

## Image Data



Western blot analysis of CD14 expression in Human tonsil cell lysate.

**Product Name: CD14 (16H16) Rabbit Monoclonal  
Antibody**  
**Catalog #: AMRe08209**

---



**Note**

For research use only.