

Product Name: FEN1 (9F12) Rabbit Monoclonal Antibody
Catalog #: AMRe10900

Summary

Production Name	FEN1 (9F12) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
Purification	Affinity purification

Immunogen

Gene Name	FEN1 {ECO:0000255 HAMAP-Rule:MF_03140}
Alternative Names	DNase IV; FEN1; Flap endonuclease 1; Flap structure specific endonuclease 1; hFEN-1; hFEN1; Maturation factor 1; MF1; Rad2;
Gene ID	2237.0
SwissProt ID	P39748.A synthetic peptide of human FEN1

Application

Dilution Ratio	WB: 1:1000
Molecular Weight	43kDa


Product Name: FEN1 (9F12) Rabbit Monoclonal Antibody
Catalog #: AMRe10900

Background

Acts as a genome stabilization factor that prevents flaps from equilibrating into structures that lead to duplications and deletions. Also possesses 5'-3' exonuclease activity on nicked or gapped double-stranded DNA, and exhibits RNase H activity. Also involved in replication and repair of rDNA and in repairing mitochondrial DNA. Structure-specific nuclease with 5'-flap endonuclease and 5'-3' exonuclease activities involved in DNA replication and repair. During DNA replication, cleaves the 5'-overhanging flap structure that is generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. It enters the flap from the 5'-end and then tracks to cleave the flap base, leaving a nick for ligation. Also involved in the long patch base excision repair (LP-BER) pathway, by cleaving within the apurinic/apyrimidinic (AP) site-terminated flap. Acts as a genome stabilization factor that prevents flaps from equilibrating into structures that lead to duplications and deletions. Also possesses 5'-3' exonuclease activity on nicked or gapped double-stranded DNA, and exhibits RNase H activity. Also involved in replication and repair of rDNA and in repairing mitochondrial DNA.

Research Area

Image Data



Western blot detection of FEN1 in HeLa, MCF-7, U251, HL-60 using FEN1 antibody (1:1000 diluted)

Note

For research use only.