

**Product Name: FOXA1 (13U2) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe11072**

---

## Summary

<b>Production Name</b>	FOXA1 (13U2) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	FOXA1
<b>Alternative Names</b>	FOXA1; Forkhead box A1; Forkhead box protein A1; HNF-3-alpha; TCF3A; HNF3A; Transcription factor 3A;
<b>Gene ID</b>	3169.0
<b>SwissProt ID</b>	P55317.Recombinant protein of human FOXA1

## Application

<b>Dilution Ratio</b>	WB: 1:1000-1:5000
<b>Molecular Weight</b>	49kDa

**Product Name: FOXA1 (13U2) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe11072**




## Background

Forkhead box protein A1 is a transcription factor required for the development of endoderm-derived organs, such as liver, lung, and prostate. FoxA1 functions as a pioneer factor that is recruited primarily to the distant enhancers to change chromatin structure for transcription in a cell type-specific manner. Involved in regulation of apoptosis by inhibiting the expression of BCL2. Involved in cell cycle regulation by activating expression of CDKN1B, alone or in conjunction with BRCA1. Transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. Binds DNA with the consensus sequence 5'- [AC]A[AT]T[AG]TT[GT][AG][CT]T[CT]-3' (By similarity). Proposed to play a role in translating the epigenetic signatures into cell type-specific enhancer-driven transcriptional programs. Its differential recruitment to chromatin is dependent on distribution of histone H3 methylated at 'Lys-5' (H3K4me2) in estrogen-regulated genes. Involved in the development of multiple endoderm-derived organ systems such as liver, pancreas, lung and prostate; FOXA1 and FOXA2 seem to have at least in part redundant roles (By similarity). Modulates the transcriptional activity of nuclear hormone receptors. Is involved in ESR1-mediated transcription; required for ESR1 binding to the NKX2-1 promoter in breast cancer cells; binds to the RPRM promoter and is required for the estrogen-induced repression of RPRM. Involved in regulation of apoptosis by inhibiting the expression of BCL2. Involved in cell cycle regulation by activating expression of CDKN1B, alone or in conjunction with BRCA1. Originally described as a transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis.

## Research Area

## Image Data



Western blot detection of FOXA1 in C6 cell lysates using FOXA1 antibody(1:1000 diluted).

**Product Name: FOXA1 (13U2) Rabbit Monoclonal  
Antibody**  
**Catalog #: AMRe11072**



---

**Note**

For research use only.