



Catalog #: AMRe12339



# **Summary**

ICAM3 / CD50 (11F8) Rabbit Monoclonal Antibody **Production Name** 

Description Rabbit Monoclonal Antibody

Host Rabbit **Application** WB.ELISA Reactivity Human

### **Performance**

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

# **Immunogen**

**Gene Name** ICAM3

**Alternative Names** CD50; CDw50; ICAM 3; ICAM3; ICAMR; Intercellular adhesion molecule 3;

Gene ID 3385.0 SwissProt ID P32942.

# **Application**

**Dilution Ratio** WB 1:500-1:2000

**Molecular Weight** 60kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Catalog #: AMRe12339




### **Background**

ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). ICAM3 is also a ligand for integrin alpha-D/beta-2. ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2) (PubMed: <a href="http://www.uniprot.org/citations/1448173" target=" blank">1448173</a>). ICAM3 is also a ligand for integrin alpha-D/beta-2. In association with integrin alpha- L/beta-2, contributes to apoptotic neutrophil phagocytosis by macrophages (PubMed: <a href="http://www.uniprot.org/citations/23775590" target="blank">23775590</a>).

#### **Research Area**

## **Image Data**



Western blot analysis of ICAM3 / CD50 expression in Ramos cell lysate.

#### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838