

**Product Name: MCM7 (3E12) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe13727**

---

## Summary

|                        |  |
|------------------------|--|
| <b>Production Name</b> | MCM7 (3E12) Rabbit Monoclonal Antibody |
| <b>Description</b>     | Rabbit Monoclonal Antibody             |
| <b>Host</b>            | Rabbit                                 |
| <b>Application</b>     | WB                                     |
| <b>Reactivity</b>      | Human                                  |

## Performance

|                     |  |
|---------------------|--|
| <b>Conjugation</b>  | Unconjugated   |
| <b>Modification</b> | Unmodified   |
| <b>Isotype</b>      | IgG  |
| <b>Clonality</b>    | Monoclonal   |
| <b>Form</b>         | Liquid   |
| <b>Storage</b>      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.                     |
| <b>Buffer</b>       | Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA. |
| <b>Purification</b> | Affinity purification  |

## Immunogen

|                          |  |
|--------------------------|--|
| <b>Gene Name</b>         | MCM7   |
| <b>Alternative Names</b> | CDC47; MCM2; Mcm7; Minichromosome Maintenance 7; P1CDC47; P85MCM; PNAS146; |
| <b>Gene ID</b>           | 4176.0   |
| <b>SwissProt ID</b>      | P33993.A synthetic peptide of human MCM7                                   |

## Application

|                         |            |
|-------------------------|------------|
| <b>Dilution Ratio</b>   | WB: 1:1000 |
| <b>Molecular Weight</b> | 81kDa      |

**Product Name: MCM7 (3E12) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe13727**




## Background

Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for S-phase checkpoint activation upon UV-induced damage.

## Research Area

## Image Data



Western blot detection of MCM7 in K562,C6,3T3,HeLa cell lysates using MCM7 antibody(1:1000 diluted).

## Note

For research use only.