

Product Name: OPRD1 (18U12) Rabbit Monoclonal Antibody
Catalog #: AMRe15359



Summary

Production Name	OPRD1 (18U12) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	OPRD1
Alternative Names	Delta type opioid receptor; Delta type opioid receptor DOR1; DOR 1; mDOR; Nbor; Opioid receptor delta 1; OPRD 1;
Gene ID	4985.0
SwissProt ID	P41143.

Application

Dilution Ratio	WB 1:1000~1:2000
Molecular Weight	40kDa

Product Name: OPRD1 (18U12) Rabbit Monoclonal Antibody
Catalog #: AMRe15359




Background

The opioid receptors are G-protein coupled, seven-transmembrane domain receptors for enkephalins, endorphins, and dynorphins. Signaling leads to the inhibition of adenylate cyclase activity. Inhibits neurotransmitter release by reducing calcium ion currents and increasing potassium ion conductance. Plays a role in the perception of pain and in opiate-mediated analgesia. Plays a role in developing analgesic tolerance to morphine. G-protein coupled receptor that functions as receptor for endogenous enkephalins and for a subset of other opioids. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Signaling leads to the inhibition of adenylate cyclase activity. Inhibits neurotransmitter release by reducing calcium ion currents and increasing potassium ion conductance. Plays a role in the perception of pain and in opiate-mediated analgesia. Plays a role in developing analgesic tolerance to morphine.

Research Area

Image Data



Western blot analysis of OPRD1 expression in SH-SY5Y cell lysate.

Note

For research use only.