

**Product Name: Pleiotrophin (12Z12) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe16264**

---

## Summary

<b>Production Name</b>	Pleiotrophin (12Z12) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	PTN
<b>Alternative Names</b>	HARP; HBBM; HBGAM; HBGF8; HBNF; HBNF1; Heparin binding growth factor 8; NEGF1; Neurite growth promoting factor 1; OSF1; Pleiotrophin; PTN;
<b>Gene ID</b>	5764.0
<b>SwissProt ID</b>	P21246.A synthetic peptide of human PTN

## Application

<b>Dilution Ratio</b>	WB: 1:1000-1:5000
<b>Molecular Weight</b>	19kDa

**Product Name: Pleiotrophin (12Z12) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe16264**

---

## Background

Heparin binding mitogenic protein. Has neurite extension activity. Secreted growth factor that mediates its signal through cell- surface proteoglycan and non-proteoglycan receptors (PubMed:<a href="http://www.uniprot.org/citations/16814777" target="\_blank">16814777</a>, PubMed:<a href="http://www.uniprot.org/citations/11278720" target="\_blank">11278720</a>, PubMed:<a href="http://www.uniprot.org/citations/19141530" target="\_blank">19141530</a>). Binds cell-surface proteoglycan receptor via their chondroitin sulfate (CS) groups (PubMed:<a href="http://www.uniprot.org/citations/26896299" target="\_blank">26896299</a>, PubMed:<a href="http://www.uniprot.org/citations/27445335" target="\_blank">27445335</a>). Thereby regulates many processes like cell proliferation, cell survival, cell growth, cell differentiation and cell migration in several tissues namely neuron and bone (PubMed:<a href="http://www.uniprot.org/citations/1733956" target="\_blank">1733956</a>, PubMed:<a href="http://www.uniprot.org/citations/1768439" target="\_blank">1768439</a>, PubMed:<a href="http://www.uniprot.org/citations/11278720" target="\_blank">11278720</a>, PubMed:<a href="http://www.uniprot.org/citations/19141530" target="\_blank">19141530</a>, PubMed:<a href="http://www.uniprot.org/citations/27445335" target="\_blank">27445335</a>, PubMed:<a href="http://www.uniprot.org/citations/30667096" target="\_blank">30667096</a>, PubMed:<a href="http://www.uniprot.org/citations/19442624" target="\_blank">19442624</a>). Also plays a role in synaptic plasticity and learning-related behavior by inhibiting long-term synaptic potentiation (By similarity). Binds PTPRZ1, leading to neutralization of the negative charges of the CS chains of PTPRZ1, inducing PTPRZ1 clustering, thereby causing the dimerization and inactivation of its phosphatase activity leading to increased tyrosine phosphorylation of each of the PTPRZ1 substrates like ALK, CTNNB1 or AFAP1L2 in order to activate the PI3K-AKT pathway (PubMed:<a href="http://www.uniprot.org/citations/17681947" target="\_blank">17681947</a>, PubMed:<a href="http://www.uniprot.org/citations/27445335" target="\_blank">27445335</a>, PubMed:<a href="http://www.uniprot.org/citations/30667096" target="\_blank">30667096</a>, PubMed:<a href="http://www.uniprot.org/citations/16814777" target="\_blank">16814777</a>, PubMed:<a href="http://www.uniprot.org/citations/10706604" target="\_blank">10706604</a>). Through PTPRZ1 binding controls oligodendrocyte precursor cell differentiation by enhancing the phosphorylation of AFAP1L2 in order to activate the PI3K-AKT pathway (PubMed:<a href="http://www.uniprot.org/citations/27445335" target="\_blank">27445335</a>, PubMed:<a href="http://www.uniprot.org/citations/30667096" target="\_blank">30667096</a>). Forms a complex with PTPRZ1 and integrin alpha-V/beta-3 (ITGAV:ITGB3) that stimulates endothelial cell migration through SRC dephosphorylation and activation that consequently leads to ITGB3 'Tyr-773' phosphorylation (PubMed:<a href="http://www.uniprot.org/citations/19141530" target="\_blank">19141530</a>). In adult hippocampus promotes dendritic arborization, spine development, and functional integration and connectivity of newborn granule neurons through ALK by activating AKT signaling pathway (By similarity). Binds GPC2 and chondroitin sulfate proteoglycans (CSPGs) at the neuron surface, leading to abrogation of binding between PTPRS and CSPGs and neurite outgrowth promotion (By similarity). Binds SDC3 and mediates bone formation by recruiting and attaching osteoblasts/osteoblast precursors to the


**Product Name: Pleiotrophin (12Z12) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe16264**

---

sites for new bone deposition (By similarity). Binds ALK and promotes cell survival and cell proliferation through MAPK pathway activation (PubMed:<a href="http://www.uniprot.org/citations/11278720" target="\_blank">11278720</a>). Inhibits proliferation and enhances differentiation of neural stem cells by inhibiting FGF2-induced fibroblast growth factor receptor signaling pathway (By similarity). Mediates regulatory mechanisms in normal hemostasis and in hematopoietic regeneration and in maintaining the balance of myeloid and lymphoid regeneration (By similarity). In addition may play a role in the female reproductive system, auditory response and the progesterone-induced decidualization pathway (By similarity).

## Research Area

## Image Data



Western blot detection of PTN in C6,Rat Brain cell lysates using PTN antibody(1:1000 diluted).

## Note

For research use only.