

Product Name: RPE65 (5T1) Rabbit Monoclonal Antibody
Catalog #: AMRe17356

Summary

Production Name	RPE65 (5T1) Rabbit Monoclonal Antibody
Description	Rabbit Monoclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Monoclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Purification	Affinity purification

Immunogen

Gene Name	RPE65
Alternative Names	LCA2; mRPE65; p63; rd12; Retinal pigment epithelium specific 61 kDa protein; RP20; RPE65; sRPE65;
Gene ID	6121.0
SwissProt ID	Q16518.

Application

Dilution Ratio	WB 1:500-1:2000
Molecular Weight	61kDa


Product Name: RPE65 (5T1) Rabbit Monoclonal Antibody
Catalog #: AMRe17356

Background

Plays important roles in the production of 11-cis retinal and in visual pigment regeneration. The soluble form binds vitamin A (all-trans-retinol), making it available for LRAT processing to all-trans-retinyl ester. The membrane form, palmitoylated by LRAT, binds all-trans-retinyl esters, making them available for IMH (isomerohydrolase) processing to all-cis-retinol. Critical isomerohydrolase in the retinoid cycle involved in regeneration of 11-cis-retinal, the chromophore of rod and cone opsins. Catalyzes the cleavage and isomerization of all-trans-retinyl fatty acid esters to 11-cis-retinol which is further oxidized by 11-cis retinol dehydrogenase to 11-cis-retinal for use as visual chromophore (PubMed:16116091). Essential for the production of 11-cis retinal for both rod and cone photoreceptors (PubMed:17848510). Also capable of catalyzing the isomerization of lutein to meso-zeaxanthin an eye- specific carotenoid (PubMed:28874556). The soluble form binds vitamin A (all-trans-retinol), making it available for LRAT processing to all- trans-retinyl ester. The membrane form, palmitoylated by LRAT, binds all-trans-retinyl esters, making them available for IMH (isomerohydrolase) processing to all-cis-retinol. The soluble form is regenerated by transferring its palmitoyl groups onto 11-cis-retinol, a reaction catalyzed by LRAT (By similarity).

Research Area

Image Data



Western blot analysis of RPE65 expression in mouse eyeball lysate.

Note

For research use only.