

**Product Name: SPOP (2B14) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe18206**

---

## Summary

<b>Production Name</b>	SPOP (2B14) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB
<b>Reactivity</b>	Human

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% BSA.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	SPOP
<b>Alternative Names</b>	BTBD32; SPOP; TEF2;
<b>Gene ID</b>	8405.0
<b>SwissProt ID</b>	O43791.A synthetic peptide of human SPOP

## Application

<b>Dilution Ratio</b>	WB: 1:1000
<b>Molecular Weight</b>	42kDa

## Background

---


**Product Name: SPOP (2B14) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe18206**

---

Inhibits IPF1/PDX1 transactivation of established target promoters, such as insulin, may be by recruiting a repressor complex (By similarity). In complex with CUL3, involved in ubiquitination of BMI1, H2AFY and DAXX, and probably also in ubiquitination and proteasomal degradation of Gli2 or Gli3. Component of a cullin-RING-based BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex that mediates the ubiquitination of target proteins, leading most often to their proteasomal degradation. In complex with CUL3, involved in ubiquitination and proteasomal degradation of BRMS1, DAXX, PDX1/IPF1, GLI2 and GLI3. In complex with CUL3, involved in ubiquitination of MACROH2A1 and BMI1; this does not lead to their proteasomal degradation. Inhibits transcriptional activation of PDX1/IPF1 targets, such as insulin, by promoting PDX1/IPF1 degradation. The cullin-RING-based BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex containing homodimeric SPOP has higher ubiquitin ligase activity than the complex that contains the heterodimer formed by SPOP and SPOPL. Involved in the regulation of bromodomain and extra-terminal motif (BET) proteins BRD2, BRD3, BRD4 stability (PubMed:<a href="http://www.uniprot.org/citations/32109420" target="\_blank">32109420</a>).

## Research Area

## Image Data



Western blot detection of SPOP in MCF-7 using SPOP antibody.

## Note

For research use only.