

**Product Name: UBA52 (9P19) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe19508**

---

## Summary

<b>Production Name</b>	UBA52 (9P19) Rabbit Monoclonal Antibody
<b>Description</b>	Rabbit Monoclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	UBA52
<b>Alternative Names</b>	60S ribosomal protein L40; CEP52; HUBCEP52; RPL40; UBA 52; Ubiquitin 52 amino acid fusion protein; Ubiquitin 60S ribosomal protein L40; Ubiquitin carboxyl extension protein 52; Ubiquitin CEP52;
<b>Gene ID</b>	7311.0
<b>SwissProt ID</b>	P62987.

## Application

<b>Dilution Ratio</b>	WB 1:500-1:2000
<b>Molecular Weight</b>	15kDa

---

**Product Name: UBA52 (9P19) Rabbit Monoclonal Antibody**  
**Catalog #: AMRe19508**

---




## Background

UBA52 is a fusion protein consisting of ubiquitin at the N terminus and ribosomal protein L40 at the C terminus, a C-terminal extension protein (CEP). Multiple processed pseudogenes derived from this gene are present in the genome. [Ubiquitin]: Exists either covalently attached to another protein, or free (unanchored). When covalently bound, it is conjugated to target proteins via an isopeptide bond either as a monomer (monoubiquitin), a polymer linked via different Lys residues of the ubiquitin (polyubiquitin chains) or a linear polymer linked via the initiator Met of the ubiquitin (linear polyubiquitin chains). Polyubiquitin chains, when attached to a target protein, have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-6-linked may be involved in DNA repair; Lys-11-linked is involved in ERAD (endoplasmic reticulum-associated degradation) and in cell cycle regulation; Lys-29-linked is involved in lysosomal degradation; Lys-33-linked is involved in kinase modification; Lys-48-linked is involved in protein degradation via the proteasome; Lys-63-linked is involved in endocytosis, DNA-damage responses as well as in signaling processes leading to activation of the transcription factor NF-kappa-B. Linear polymer chains formed via attachment by the initiator Met lead to cell signaling. Ubiquitin is usually conjugated to Lys residues of target proteins, however, in rare cases, conjugation to Cys or Ser residues has been observed. When polyubiquitin is free (unanchored- polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling.

## Research Area

## Image Data



Western blot analysis of UBA52 expression in 293T cell lysate.

## Note

For research use only.

**Product Name: UBA52 (9P19) Rabbit Monoclonal  
Antibody**  
**Catalog #: AMRe19508**

---

