



Catalog #: APRab03382



Summary

Phospho-Chk2 (Thr68) Rabbit Polyclonal Antibody **Production Name**

Description Primary antibody

Host Rabbit

Application WB,IHC-P,ICC/IF Reactivity Human, Mouse, Rat

Performance

Conjugation Unconjugated Modification Phosphorylated

Isotype IgG

Clonality Polyclonal Antibody

Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Purification Affinity Purified

Immunogen

Gene Name CHEK2

CHEK2; CDS1; CHK2; RAD53; Serine/threonine-protein kinase Chk2; CHK2 checkpoint

Alternative Names homolog; Cds1 homolog; Hucds1; hCds1; Checkpoint kinase 2

Gene ID 11200 **SwissProt ID** O96017

Application

Dilution Ratio WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200

Molecular Weight Calculated MW: 61 kDa; Observed MW: 61 kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Catalog #: APRab03382




Background

These are known to be preferred sites for phosphorylation by ATM/ATR kinases. After DNA damage by ionizing radiation (IR), UV irradiation, or hydroxyurea treatment, Thr68 and other sites in this region become phosphorylated by ATM/ATR. The SQ/TQ cluster domain, therefore, seems to have a regulatory function.

Research Area

Epigenetics and Nuclear Signaling

Image Data



Western blot analysis of Phospho-Chk2(Thr68) in Jurkat treated with UV lysates using Phospho- Chk2 (Thr68) antibody.

Note

For research use only.