## Product Name: ALK (phospho Tyr1507) Rabbit

Polyclonal Antibody Catalog #: APRab04221



#### **Summary**

**Production Name** ALK (phospho Tyr1507) Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

**Application** ELISA,IHC,WB

**Reactivity** Human, Mouse, Monkey

#### **Performance**

**Conjugation** Unconjugated

**Modification** Phospho Antibody

**Isotype** IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

#### **Immunogen**

Gene Name ALK

Alternative Names ALK; ALK tyrosine kinase receptor; Anaplastic lymphoma kinase; CD antigen CD246

**Gene ID** 238.0

Q9UM73.The antiserum was produced against synthesized peptide derived from **SwissProt ID** 

human ALK around the phosphorylation site of Tyr1507. AA range:1473-1522

### **Application**

**Dilution Ratio** WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000..

Molecular Weight 150-240kD

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: ALK (phospho Tyr1507) Rabbit

Polyclonal Antibody Catalog #: APRab04221




#### **Background**

This gene encodes a receptor tyrosine kinase, which belongs to the insulin receptor superfamily. This protein comprises an extracellular domain, an hydrophobic stretch corresponding to a single pass transmembrane region, and an intracellular kinase domain. It plays an important role in the development of the brain and exerts its effects on specific neurons in the nervous system. This gene has been found to be rearranged, mutated, or amplified in a series of tumours including anaplastic large cell lymphomas, neuroblastoma, and non-small cell lung cancer. The chromosomal rearrangements are the most common genetic alterations in this gene, which result in creation of multiple fusion genes in tumourigenesis, including ALK (chromosome 2)/EML4 (chromosome 2), ALK/RANBP2 (chromosome 2), ALK/ATIC (chromosome 2), ALK/TFG (chromosome 3), ALK/NPM1 (chromosome 5), ALK/SQSTM1 (chromosomecatalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate, disease: A chromosomal aberration involving ALK is associated with anaplastic large-cell lymphoma (ALCL). Translocation t(2;17)(p23;q25) with ALO17., disease: A chromosomal aberration involving ALK is associated with inflammatory myofibroblastic tumors (IMTs). Translocation t(2;11)(p23;p15) with CARS; translocation t(2;4) (p23;q21) with SEC31A, disease: A chromosomal aberration involving ALK is found in a form of non-Hodgkin lymphoma. Translocation t(2;5)(p23;q35) with NPM1. The resulting chimeric NPM1-ALK protein homodimerize and the kinase becomes constitutively activated. The constitutively active fusion proteins are responsible for 5-10% of non-Hodgkin lymphomas, function: Orphan receptor with a tyrosine-protein kinase activity. Appears to play an important role in the normal development and function of the nervous system. Phosphorylates almost exclusively at the first tyrosine of the Y-xx-x-Y-Y motif., PTM:N-glycosylated, similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily,,similarity:Contains 1 LDL-receptor class A domain,,similarity:Contains 1 protein kinase domain, similarity: Contains 2 MAM domains, subunit: Homodimer. When bound to ligand, tissue specificity: Expressed in brain and CNS. Also expressed in the small intestine and testis, but not in normal lymphoid cells.,

#### **Research Area**

#### **Image Data**



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# Product Name: ALK (phospho Tyr1507) Rabbit

Polyclonal Antibody Catalog #: APRab04221



Phosphopeptide (Phospho-right), using ALK (Phospho-Tyr1507) Antibody



Immunohistochemistry analysis of paraffin-embedded human brain, using ALK (Phospho-Tyr1507) Antibody. The picture on the right is blocked with the phospho peptide.

Western blot analysis of lysates from COS7 cells treated with anisomycin 25ug/ml 30 ', using ALK (Phospho-Tyr1507)

Antibody. The lane on the right is blocked with the phospho peptide.

#### Note

For research use only.