

# Summary

Production Name	CaSR (phospho Thr888) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	ELISA,IF,WB
Reactivity	Human, Mouse, Rat

### Performance

Conjugation	Unconjugated	
Modification	Phospho Antibody	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

### Immunogen

Gene Name	CASR
Alternative Names	CASR; GPRC2A; PCAR1; Extracellular calcium-sensing receptor; CaSR; Parathyroid cell
	calcium-sensing receptor; PCaR1
Gene ID	846.0
	P41180.The antiserum was produced against synthesized peptide derived from human
SwissProt ID	Calcium Sensing Receptor around the phosphorylation site of Thr888. AA range:854-
	903

# Application

Dilution Ratio	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:40000. Not yet tested in other
----------------	---

## Product Name: CaSR (phospho Thr888) Rabbit Polyclonal Antibody Catalog #: APRab04376



applications.

**Molecular Weight** 

140kD


## Background

The protein encoded by this gene is a G protein-coupled receptor that is expressed in the parathyroid hormone (PTH)producing chief cells of the parathyroid gland, and the cells lining the kidney tubule. It senses small changes in circulating calcium concentration and couples this information to intracellular signaling pathways that modify PTH secretion or renal cation handling, thus this protein plays an essential role in maintaining mineral ion homeostasis. Mutations in this gene cause familial hypocalciuric hypercalcemia, familial, isolated hypoparathyroidism, and neonatal severe primary hyperparathyroidism. [provided by RefSeq, Jul 2008], disease: Defects in CASR are the cause of autosomal dominant hypoparathyroidism (FIH) [MIM:146200]. FIH is characterized by hypocalcemia and hyperphosphatemia due to inadequate secretion of parathyroid hormone. Symptoms are seizures, tetany and cramps.,disease:Defects in CASR are the cause of familial hypocalciuric hypercalcemia type 1 (FHH) [MIM:145980]; in which the receptor has reduced activity. FHH is characterized by altered calcium homeostasis. Affected individuals exhibit mild or modest hypercalcemia, relative hypocalciuria, and inappropriately normal PTH levels., disease: Defects in CASR are the cause of neonatal severe primary hyperparathyroidism (NSHPT) [MIM:239200]; in which the receptor has reduced activity. NSHPT is a rare autosomal recessive life-threatening disorder characterized by very high serum calcium concentrations, skeletal demineralization, and parathyroid hyperplasia. In some instances NSHPT has been demonstrated to be the homozygous form of FHH., function: Senses changes in the extracellular concentration of calcium ions. The activity of this receptor is mediated by a G-protein that activates a phosphatidylinositol-calcium second messenger system, PTM:Nglycosylated., PTM: Ubiguitinated by RNF19A; which induces proteasomal degradation., similarity: Belongs to the G-protein coupled receptor 3 family., subunit: Interacts with VCP and RNF19A., tissue specificity: Found in kidney, but not in brain, lung, liver, heart, skeletal muscle, or placenta.,


#### **Research Area**

**Image Data** 






Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Calcium Sensing Receptor (Phospho-Thr888) Antibody



Immunofluorescence analysis of HeLa cells, using Calcium Sensing Receptor (Phospho-Thr888) Antibody. The picture on



Western blot analysis of lysates from LOVO cells, using Calcium Sensing Receptor (Phospho-Thr888) Antibody. The lane on the right is blocked with the phospho peptide.

#### Note

For research use only.