Polyclonal Antibody Catalog #: APRab04589



Summary

Production Name EGFR (phospho Thr693) Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application ELISA,IF,IHC,WB **Reactivity** Human,Mouse,Rat

Performance

Conjugation Unconjugated

Modification Phospho Antibody

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name EGFR

EGFR; ERBB; ERBB1; HER1; Epidermal growth factor receptor; Proto-oncogene c-ErbB-1; Alternative Names

Receptor tyrosine-protein kinase erbB-1

Gene ID 1956.0

P00533.The antiserum was produced against synthesized peptide derived from human **SwissProt ID**

EGFR around the phosphorylation site of Thr693. AA range:661-710

Application

Dilution Ratio

WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:40000. Not yet tested in

other application

 $other\ applications.$

Polyclonal Antibody Catalog #: APRab04589



Molecular Weight

175kD

Background

The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this gene are associated with lung cancer. [provided by RefSeq, Jun 2016], catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., disease:Defects in EGFR are associated with lung cancer [MIM:211980], function: Isoform 2/truncated isoform may act as an antagonist., function: Receptor for EGF, but also for other members of the EGF family, as TGF-alpha, amphiregulin, betacellulin, heparin-binding EGF-like growth factor, GP30 and vaccinia virus growth factor. Is involved in the control of cell growth and differentiation. Phosphorylates MUC1 in breast cancer cells and increases the interaction of MUC1 with C-SRC and CTNNB1/beta-catenin., miscellaneous: Binding of EGF to the receptor leads to dimerization, internalization of the EGFreceptor complex, induction of the tyrosine kinase activity, stimulation of cell DNA synthesis, and cell proliferation, online information:EGFR entry,PTM:Monoubiquitinated and polyubiquitinated upon EGF stimulation; which does not affect tyrosine kinase activity or signaling capacity but may play a role in lysosomal targeting. Polyubiquitin linkage is mainly through 'Lys-63', but linkage through 'Lys-48', 'Lys-11' and 'Lys-29' also occur., PTM: Phosphorylation of Ser-695 is partial and occurs only if Thr-693 is phosphorylated., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. EGF receptor subfamily,, similarity: Contains 1 protein kinase domain., subunit: Binds RIPK1. CBL interacts with the autophosphorylated Cterminal tail of the EGF receptor. Part of a complex with ERBB2 and either PIK3C2A or PIK3C2B. The autophosphorylated form interacts with PIK3C2B, maybe indirectly. Interacts with PELP1. Binds MUC1.,tissue specificity:Ubiquitously expressed. Isoform 2 is also expressed in ovarian cancers.,

Research Area


MAPK_ERK_Growth;MAPK_G_Protein;ErbB_HER;Calcium;Cytokine-cytokine receptor interaction;Endocytosis;Dorso-ventral axis formation;Focal adhesion;Adherens_Junction;Gap junction;Regulates Actin and Cytoskeleton;GnRH;Epithelial cell signaling in Helicobacter pylori infection;Pathways in cancer;Colorectal cancer;Pancreatic cancer;Endometrial cancer;Glioma;Prostate cancer;Melanoma;Bladder cancer;Non-small cell lung cancer;

Image Data


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Polyclonal Antibody Catalog #: APRab04589






Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right) , using EGFR (Phospho-Thr693) Antibody



Immunofluorescence analysis of HUVEC cells, using EGFR (Phospho-Thr693) Antibody. The picture on the right is blocked with the phospho peptide.




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using EGFR (Phospho-Thr693) Antibody.

The picture on the right is blocked with the phospho peptide.


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Polyclonal Antibody Catalog #: APRab04589





Western blot analysis of lysates from A431 cells, using EGFR (Phospho-Thr693) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-EGFR (T693) Polyclonal Antibody

Note

For research use only.