

Summary

Production Name	ΙκΒ-β (phospho Ser23) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	ELISA,IF,IHC,WB
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated	
Modification	Phospho Antibody	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

Immunogen

Gene Name	NFKBIB
Alternative Names	NFKBIB; IKBB; TRIP9; NF-kappa-B inhibitor beta; NF-kappa-BIB; I-kappa-B-beta; IkB-B;
	IkB-beta; IkappaBbeta; Thyroid receptor-interacting protein 9; TR-interacting protein 9;
	TRIP-9
Gene ID	4793.0
SwissProt ID	Q15653.The antiserum was produced against synthesized peptide derived from human
	IkappaB-beta around the phosphorylation site of Ser23. AA range:8-57

Application

Dilution Ratio	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:40000. Not yet tested in
----------------	--



other applications.

Molecular Weight

37kD


Background

The protein encoded by this gene belongs to the NF-kappa-B inhibitor family, which inhibit NF-kappa-B by complexing with, and trapping it in the cytoplasm. Phosphorylation of serine residues on these proteins by kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B, which translocates to the nucleus to function as a transcription factor. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2011],function:Inhibits NF-kappa-B by complexing with and trapping it in the cytoplasm. However, the unphosphorylated form resynthesized after cell stimulation is able to bind NF-kappa-B allowing its transport to the nucleus and protecting it to further IKBA-dependent inactivation. Association with inhibitor kappa B-interacting NKIRAS1 and NKIRAS2 prevent its phosphorylated; followed by degradation. Interaction with NKIRAS1 and NKIRAS2 probably prevents phosphorylated; followed by degradation. Interaction with NKIRAS1 and NKIRAS2 probably prevents with THRB (via ligand-binding domain). Interacts with RELA and REL. Interacts with COMMD1 and inhibitor kappa B-interacting Ras-like NKIRAS1 and NKIRAS2, tissue specificity:Expressed in all tissues examined.,


Research Area


Chemokine;NOD-like receptor;RIG-I-like receptor;Cytosolic DNA-sensing pathway;T_Cell_Receptor;B_Cell_Antigen;Neurotrophin;Adipocytokine;

Image Data




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using IkappaB-beta (Phospho-Ser23) Antibody







Immunofluorescence analysis of HeLa cells treated with TNF-a 20nM 15 ', using IkappaB-beta (Phospho-Ser23) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using IkappaB-beta (Phospho-Ser23) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HeLa cells treated with TNF-a 20ng/ml 5 ', using IkappaB-beta (Phospho-Ser23) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-I κ B- β (S23) Polyclonal Antibody

Note

For research use only.