

Product Name: MEK Kinase-1 (phospho Thr1402) Rabbit Polyclonal Antibody
Catalog #: APRab04999

Summary

| | |
|------------------------|---|
| Production Name | MEK Kinase-1 (phospho Thr1402) Rabbit Polyclonal Antibody |
| Description | Rabbit Polyclonal Antibody |
| Host | Rabbit |
| Application | ELISA,IHC,WB, |
| Reactivity | Human,Mouse,Rat |

Performance

| | |
|---------------------|--|
| Conjugation | Unconjugated |
| Modification | Phospho Antibody |
| Isotype | IgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification |

Immunogen

| | |
|--------------------------|--|
| Gene Name | MAP3K1 |
| Alternative Names | MAP3K1; MAPKKK1; MEKK; MEKK1; Mitogen-activated protein kinase kinase kinase 1; MAPK/ERK kinase kinase 1; MEK kinase 1; MEKK 1 |
| Gene ID | 4214.0 |
| SwissProt ID | Q13233.The antiserum was produced against synthesized peptide derived from human MAP3K1 around the phosphorylation site of Thr1402. AA range:1368-1417 |

Application

| | |
|-------------------------|---|
| Dilution Ratio | WB 1:500 - 1:2000 IHC 1:100 - 1:300. ELISA: 1:40000.. |
| Molecular Weight | 160kD |

Product Name: MEK Kinase-1 (phospho Thr1402) Rabbit Polyclonal Antibody
Catalog #: AP Rab04999


Background

The protein encoded by this gene is a serine/threonine kinase and is part of some signal transduction cascades, including the ERK and JNK kinase pathways as well as the NF-kappa-B pathway. The encoded protein is activated by autophosphorylation and requires magnesium as a cofactor in phosphorylating other proteins. This protein has E3 ligase activity conferred by a plant homeodomain (PHD) in its N-terminus and phospho-kinase activity conferred by a kinase domain in its C-terminus. [provided by RefSeq, Mar 2012], catalytic activity: ATP + a protein = ADP + a phosphoprotein., cofactor: Magnesium., enzyme regulation: Activated by autophosphorylation on Thr-1400 and Thr-1412 following oligomerization., function: Component of a protein kinase signal transduction cascade. Activates the ERK and JNK kinase pathways by phosphorylation of MAP2K1 and MAP2K4. Activates CHUK and IKBKB, the central protein kinases of the NF-kappa-B pathway., PTM: Autophosphorylated., similarity: Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase kinase subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 1 RING-type zinc finger., similarity: Contains 1 SWIM-type zinc finger., subunit: Binds both upstream activators and downstream substrates in multimolecular complexes through its N-terminus. Oligomerizes after binding MAP4K2 or TRAF2. Interacts with AXIN1.,

Research Area


SAPK_JNK; Regulation of Actin Dynamics; Cell Growth; Stem cell pathway; MAPK_ERK_Growth; MAPK_G_Protein; B Cell Receptor

Image Data




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using MAP3K1 (Phospho-Thr1402) Antibody


Product Name: MEK Kinase-1 (phospho Thr1402) Rabbit Polyclonal Antibody
Catalog #: APRab04999



Immunohistochemistry analysis of paraffin-embedded human brain, using MAP3K1 (Phospho-Thr1402) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from Jurkat cells, using MAP3K1 (Phospho-Thr1402) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of NIH-3T3 cells using Phospho-MEK Kinase-1 (T1402) Polyclonal Antibody diluted at 1: 2000

Note

For research use only.