

**Product Name: Myt 1 (phospho Ser83) Rabbit Polyclonal Antibody**  
**Catalog #: APRab05061**

---

## Summary

<b>Production Name</b>	Myt 1 (phospho Ser83) Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC,ELISA
<b>Reactivity</b>	Human,Rat,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Phospho Antibody
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	PKMYT1
<b>Alternative Names</b>	PKMYT1; MYT1; Membrane-associated tyrosine- and threonine-specific cdc2-inhibitory kinase; Myt1 kinase
<b>Gene ID</b>	9088.0
<b>SwissProt ID</b>	Q99640.The antiserum was produced against synthesized peptide derived from human MYT1 around the phosphorylation site of Ser83. AA range:49-98

## Application

<b>Dilution Ratio</b>	IHC 1:100 - 1:300. ELISA: 1:5000..
<b>Molecular Weight</b>	

**Product Name: Myt 1 (phospho Ser83) Rabbit Polyclonal Antibody**  
**Catalog #: AP Rab05061**




## Background

This gene encodes a member of the serine/threonine protein kinase family. The encoded protein is a membrane-associated kinase that negatively regulates the G2/M transition of the cell cycle by phosphorylating and inactivating cyclin-dependent kinase 1. The activity of the encoded protein is regulated by polo-like kinase 1. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, May 2012], catalytic activity: ATP + a protein = ADP + a phosphoprotein., domain: The membrane-association motif is essential for the localization to membrane of Golgi stack. According to some authors, it is a transmembrane domain; the existence of a transmembrane region is however unproven., enzyme regulation: Negatively regulated by hyperphosphorylation during mitosis. The hyperphosphorylated form does not associate with CCNB1-CDC2 complexes. The PLK1 protein kinase may be required for mitotic phosphorylation., function: Acts as a negative regulator of entry into mitosis (G2 to M transition) by phosphorylation of the cdc2 kinase specifically when cdc2 is complexed to cyclins. Mediates phosphorylation of cdc2 predominantly on 'Thr-14'. Also involved in Golgi fragmentation. May be involved in phosphorylation of cdc2 on 'Tyr-15' to a lesser degree, however tyrosine kinase activity is unclear and may be indirect. May be a downstream target of Notch signaling pathway during eye development., PTM: Autophosphorylated. Phosphorylated by CDC2-CCNB1 complexes on undefined serine and threonine residues. The phosphorylation by CDC2-CCNB1 complexes may inhibit the catalytic activity., sequence caution: Chimeric cDNA., similarity: Belongs to the protein kinase superfamily., similarity: Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. WEE1 subfamily., similarity: Contains 1 protein kinase domain., subunit: Interacts with CDC2-CCNB1 complex. Can also interact with PIN1 when phosphorylated by CDC2-CCNB1.,

## Research Area

Cell\_Cycle\_G1S; Cell\_Cycle\_G2M\_DNA; Oocyte meiosis; Progesterone-mediated oocyte maturation;


## Image Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using MYT1 (Phospho-Ser83) Antibody

**Product Name: Myt 1 (phospho Ser83) Rabbit Polyclonal Antibody**  
**Catalog #: APRab05061**

---



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using MYT1 (Phospho-Ser83) Antibody.  
The picture on the right is blocked with the phospho peptide.

### **Note**

For research use only.