

**Product Name: p47-phox (phospho Ser328) Rabbit Polyclonal Antibody**  
**Catalog #: APRab05159**

---

## Summary

<b>Production Name</b>	p47-phox (phospho Ser328) Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC,ELISA
<b>Reactivity</b>	Human,Mouse,Rat,Cow

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Phospho Antibody
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	NCF1
<b>Alternative Names</b>	NCF1; NOXO2; SH3PXD1A; Neutrophil cytosol factor 1; NCF-1; 47 kDa autosomal chronic granulomatous disease protein; 47 kDa neutrophil oxidase factor; NCF-47K; Neutrophil NADPH oxidase factor 1; Nox organizer 2; Nox-organizing protein 2; SH3
<b>Gene ID</b>	653361.0
<b>SwissProt ID</b>	P14598.The antiserum was produced against synthesized peptide derived from human Neutrophil Cytosol Factor 1 around the phosphorylation site of Ser328. AA range:301-350

## Application

<b>Dilution Ratio</b>	IHC 1:100 - 1:300. ELISA: 1:10000.
-----------------------	------------------------------------

---

**Product Name: p47-phox (phospho Ser328) Rabbit Polyclonal Antibody**  
**Catalog #: APRab05159**



---

## Molecular Weight


## Background

The protein encoded by this gene is a 47 kDa cytosolic subunit of neutrophil NADPH oxidase. This oxidase is a multicomponent enzyme that is activated to produce superoxide anion. Mutations in this gene have been associated with chronic granulomatous disease. [provided by RefSeq, Jul 2008],disease:Defects in NCF1 are the cause of chronic granulomatous disease autosomal recessive cytochrome-b-positive type 1 (CGD1) [MIM:233700]. Chronic granulomatous disease is a genetically heterogeneous disorder characterized by the inability of neutrophils and phagocytes to kill microbes that they have ingested. Patients suffer from life-threatening bacterial/fungal infections.,function:NCF2, NCF1, and a membrane bound cytochrome b558 are required for activation of the latent NADPH oxidase (necessary for superoxide production),.online information:NCF1 deficiency database,similarity:Contains 1 PX (phox homology) domain,.similarity:Contains 2 SH3 domains,.subunit:Interacts with NOXA1,.

## Research Area

Chemokine;Fc gamma R-mediated phagocytosis;Leukocyte transendothelial migration;


## Image Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right) , using Neutrophil Cytosol Factor 1 (Phospho-Ser328) Antibody

**Product Name: p47-phox (phospho Ser328) Rabbit  
Polyclonal Antibody  
Catalog #: APRab05159**

---



Immunohistochemistry analysis of paraffin-embedded human tonsil, using Neutrophil Cytosol Factor 1 (Phospho-Ser328) Antibody. The picture on the right is blocked with the phospho peptide.

**Note**

For research use only.