

Antibody

Catalog #: APRab05376



Summary

RIP2 (phospho Ser176) Rabbit Polyclonal Antibody **Production Name**

Description Rabbit Polyclonal Antibody

Host Rabbit

Application ELISA,IHC,WB, Reactivity Human, Mouse

Performance

Unconjugated Conjugation

Modification Phospho Antibody

Isotype IgG

Clonality Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name RIPK2

RIPK2; CARDIAK; RICK; RIP2; Receptor-interacting serine/threonine-protein kinase 2;

Alternative Names CARD-containing interleukin-1 beta-converting enzyme-associated kinase; CARD-

containing IL-1 beta ICE-kinase; RIP-like-interacting CLARP kinase; Receptor-in

Gene ID 8767.0

O43353. The antiserum was produced against synthesized peptide derived from human SwissProt ID

RIPK2 around the phosphorylation site of Ser176. AA range:146-195

Application

Dilution Ratio WB 1:500 - 1:2000 IHC 1:100 - 1:300. ELISA: 1:40000...

Molecular Weight 70kD

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Catalog #: APRab05376




Background

This gene encodes a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases. The encoded protein contains a C-terminal caspase activation and recruitment domain (CARD), and is a component of signaling complexes in both the innate and adaptive immune pathways. It is a potent activator of NF-kappaB and inducer of apoptosis in response to various stimuli. [provided by RefSeq, Jul 2008],catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Activates pro-caspase-1 and pro-caspase-8. Potentiates CASP8-mediated apoptosis. Activates NF-kappa-B., PTM: Autophosphorylated. Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family., similarity: Contains 1 CARD domain., similarity: Contains 1 protein kinase domain., subunit: Binds to CFLAR/CLARP and CASP1 via their CARD domains. Binds to BIRC3/c-IAP1 and BIRC2/c-IAP2, TRAF1, TRAF2, TRAF5 and TRAF6. May be a component of both the TNFRSF1A and TNRFSF5/CD40 receptor complex.,tissue specificity:Detected in heart, brain, placenta, lung, peripheral blood leukocytes, spleen, kidney, testis, prostate, pancreas and lymph node.,

Research Area

NOD-like receptor; Neurotrophin;

Image Data




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using RIPK2 (Phospho-Ser176) Antibody

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




Catalog #: APRab05376






Immunohistochemistry analysis of paraffin-embedded human brain, using RIPK2 (Phospho-Ser176) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with UV 15 ', using RIPK2 (Phospho-Ser176) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of 293 cells using Phospho-RIP2 (S176) Polyclonal Antibody

Note

For research use only.