

Product Name: NCoA3 (Phospho-Ser857) Rabbit Polyclonal Antibody
Catalog #: APRab05815

Summary

Production Name	NCoA3 (Phospho-Ser857) Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Phospho Antibody
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	NCOA3 AIB1 BHLHE42 RAC3 TRAM1 Nuclear receptor coactivator 3 (NCoA-3) (EC 2.3.1.48) (ACTR) (Amplified in breast cancer 1 protein) (AIB-1) (CBP-interacting protein) (pCIP) (Class E basic helix-loop-helix protein 42) (bHLHe42) (Receptor-associated coactivator 3) (RAC-3) (Steroid receptor coactivator protein 3) (SRC-3) (Thyroid hormone receptor activator molecule 1) (TRAM-1)
Alternative Names	
Gene ID	8202.0
SwissProt ID	Q9Y6Q9.

Application

Dilution Ratio	WB 1:500-2000
Molecular Weight	157kD

**Product Name: NCoA3 (Phospho-Ser857) Rabbit
Polyclonal Antibody
Catalog #: APRab05815**


Background

alternative products: Additional isoforms seem to exist, catalytic activity: Acetyl-CoA + histone = CoA + acetylhistone., domain: Contains three Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. Motifs 1 and 2 are essential for the association with nuclear receptors, and constitute the RID domain (Receptor-interacting domain), enzyme regulation: Coactivator activity on nuclear receptors and NF-kappa-B pathways is enhanced by various hormones, and the TNF cytokine, respectively. TNF stimulation probably enhances phosphorylation, which in turn activates coactivator function. In contrast, acetylation by CREBBP apparently suppresses coactivation of target genes by disrupting its association with nuclear receptors., function: Nuclear receptor coactivator that directly binds nuclear receptors and stimulates the transcriptional activities in a hormone-dependent fashion. Plays a central role in creating a multisubunit coactivator complex, which probably acts via remodeling of chromatin. Involved in the coactivation of different nuclear receptors, such as for steroids (GR and ER), retinoids (RARs and RXRs), thyroid hormone (TRs), vitamin D3 (VDR) and prostanoids (PPARs). Displays histone acetyltransferase activity. Also involved in the coactivation of the NF-kappa-B pathway via its interaction with the NFKB1 subunit., miscellaneous: NCOA3 is frequently amplified or overexpressed in breast and ovarian cancers., polymorphism: The length of the poly-Gln region is polymorphic in the normal population., PTM: Acetylated by CREBBP. Acetylation occurs in the RID domain, and disrupts the interaction with nuclear receptors and regulates its function., PTM: Methylated by CARM1., PTM: Phosphorylated by IKK complex. Regulated its function., similarity: Belongs to the SRC/p160 nuclear receptor coactivator family., similarity: Contains 1 basic helix-loop-helix (bHLH) domain., similarity: Contains 1 PAS (PER-ARNT-SIM) domain., subcellular location: Mainly cytoplasmic and weakly nuclear. Upon TNF activation and subsequent phosphorylation, it translocates from the cytoplasm to the nucleus., subunit: Interacts with CARM1 (By similarity). Present in a complex containing NCOA2, IKKA, IKKB, IKBKG and the histone acetyltransferase protein CREBBP. Interacts with CASP8AP2, NR3C1 and PCAF. Interacts with ATAD2 and this interaction is enhanced by estradiol., tissue specificity: Widely expressed. High expression in heart, skeletal muscle, pancreas and placenta. Low expression in brain, and very low in lung, liver and kidney.,

Research Area

Image Data

**Product Name: NCoA3 (Phospho-Ser857) Rabbit
Polyclonal Antibody
Catalog #: APRab05815**



Western Blot analysis of 1 MCF-7 cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution.
Secondary antibody was diluted at 1:10000

Note

For research use only.