

Product Name: 3pK Rabbit Polyclonal Antibody
Catalog #: APRab06320

Summary

Production Name	3pK Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC, WB, ELISA
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	MAPKAPK3
Alternative Names	MAPKAPK3; MAP kinase-activated protein kinase 3; MAPK-activated protein kinase 3; MAPKAP kinase 3; MAPKAP-K3; MAPKAPK-3; MK-3; Chromosome 3p kinase; 3pK
Gene ID	7867.0
SwissProt ID	Q16644. The antiserum was produced against synthesized peptide derived from human MAPK3. AA range: 301-350

Application

Dilution Ratio	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000. Not yet tested in other applications.
Molecular Weight	42kD


Background

This gene encodes a member of the Ser/Thr protein kinase family. This kinase functions as a mitogen-activated protein kinase (MAP kinase)- activated protein kinase. MAP kinases are also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This kinase was shown to be activated by growth inducers and stress stimulation of cells. In vitro studies demonstrated that ERK, p38 MAP kinase and Jun N-terminal kinase were all able to phosphorylate and activate this kinase, which suggested the role of this kinase as an integrative element of signaling in both mitogen and stress responses. This kinase was reported to interact with, phosphorylate and repress the activity of E47, which is a basic helix-loop-helix transcription factor known to be involved in the regulation of tissue-specific gene expression and catalytic activity: ATP + a protein = ADP + a phosphoprotein., function: Modulator of polycomb-mediated repression, which can be activated either by ERK, p38 and JNK. Substrate of CSBP. In vitro, phosphorylates HSPB1, BMI1/PCGF4 and TCF3., similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family., similarity: Contains 1 protein kinase domain., subcellular location: Predominantly located in the nucleus, when activated it translocates to the cytoplasm., subunit: Interacts with TCF3 and with polycomb proteins, such as PCH2 and BMI1/PCGF4., tissue specificity: Widely expressed, with a higher expression level observed in heart and skeletal muscle. No expression in brain.,

Research Area


MAPK_ERK_Growth; MAPK_G_Protein; VEGF;

Image Data




Immunohistochemistry analysis of paraffin-embedded human skeletal muscle tissue, using MAPK3 Antibody. The picture on the right is blocked with the synthesized peptide.

Product Name: 3pK Rabbit Polyclonal Antibody
Catalog #: APRab06320



Western blot analysis of lysates from COLO205 cells, using MAPK3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using 3pK Polyclonal Antibody

Note

For research use only.