

Summary

| Production Name | A Cyclase IV Rabbit Polyclonal Antibody |
|-----------------|---|
| Description | Rabbit Polyclonal Antibody |
| Host | Rabbit |
| Application | WB,ELISA |
| Reactivity | Human, Mouse, Rat |

Performance

| Conjugation | Unconjugated |
|--------------|--|
| Modification | Unmodified |
| lsotype | lgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw |
| | cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification |

Immunogen

| Gene Name | ADCY4 |
|-------------------|--|
| Alternative Names | ADCY4; Adenylate cyclase type 4; ATP pyrophosphate-lyase 4; Adenylate cyclase type |
| | IV; Adenylyl cyclase 4 |
| Gene ID | 196883.0 |
| SwissProt ID | Q8NFM4.The antiserum was produced against synthesized peptide derived from |
| | human ADCY4. AA range:195-244 |

Application

| Dilution Ratio | WB 1:500 - 1:2000. ELISA: 1:20000 |
|------------------|-----------------------------------|
| Molecular Weight | 120kD |

Product Name: A Cyclase IV Rabbit Polyclonal Antibody


Background

This gene encodes a member of the family of adenylate cyclases, which are membrane-associated enzymes that catalyze the formation of the secondary messenger cyclic adenosine monophosphate (cAMP). Mouse studies show that adenylate cyclase 4, along with adenylate cyclases 2 and 3, is expressed in olfactory cilia, suggesting that several different adenylate cyclases may couple to olfactory receptors and that there may be multiple receptor-mediated mechanisms for the generation of cAMP signals. Alternative splicing results in transcript variants. [provided by RefSeq, Nov 2010],catalytic activity:ATP = 3',5'-cyclic AMP + diphosphate.,cofactor:Binds 2 magnesium ions per subunit.,enzyme regulation:Insensitive to calcium/calmodulin. Stimulated by the G protein beta and gamma subunit complex.,function:This is a membrane-bound, calmodulin-insensitive adenylyl cyclase.,similarity:Belongs to the adenylyl cyclase class-4/guanylyl cyclase family.,similarity:Contains 2 guanylate cyclase domains.,


Research Area

Purine metabolism;Calcium;Chemokine;Oocyte meiosis;Vascular smooth muscle contraction;Gap junction;Taste transduction;GnRH;Progesterone-mediated oocyte maturation;Melanogenesis;Dilated cardiomyopathy;

Image Data



Western blot analysis of lysates from Jurkat cells, using ADCY4 Antibody. The lane on the right is blocked with the



Western Blot analysis of Jurkat cells using A Cyclase IV Polyclonal Antibody



Note For research use only.