

**Product Name: ANKT Rabbit Polyclonal Antibody**  
**Catalog #: APRab06907**



## Summary

<b>Production Name</b>	ANKT Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Rat,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	NUSAP1
<b>Alternative Names</b>	NUSAP1; ANKT; BM-037; Nucleolar and spindle-associated protein 1; NuSAP
<b>Gene ID</b>	51203.0
<b>SwissProt ID</b>	Q9BXS6.The antiserum was produced against synthesized peptide derived from human NUSAP1. AA range:392-441

## Application

<b>Dilution Ratio</b>	WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.
<b>Molecular Weight</b>	49kD

## Background


**Product Name: ANKT Rabbit Polyclonal Antibody**  
**Catalog #: APRab06907**




NUSAP1 is a nucleolar-spindle-associated protein that plays a role in spindle microtubule organization (Raemaekers et al., 2003 [PubMed 12963707]).[supplied by OMIM, Jun 2009],domain:The KEN box is required for the FZR1-dependent degradation of this protein subsequent to ubiquitination.,function:Microtubule-associated protein with the capacity to bundle and stabilize microtubules (By similarity). May associate with chromosomes and promote the organization of mitotic spindle microtubules around them.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,PTM:Ubiquitinated. Ubiquitination by FZR1 may lead to proteasome-dependent degradation of this protein.,similarity:Belongs to the NUSAP family.,subcellular location:Found in the cytoplasm and nucleolus during interphase and redistributes to the mitotic spindle in prometaphase (By similarity). Localizes to the mitotic spindle during anaphase and telophase then disappears from around the chromosomes during cytokinesis (By similarity). Localizes to multiple distinct regions of chromosomes throughout mitosis.,subunit:Interacts with DNA and microtubules. Microtubule bundling is inhibited by IPO7, KPNA2 and KPNB1 while association with DNA is also inhibited by IPO7 and KPNA2.,

## Research Area

## Image Data




Western blot analysis of lysates from HT-29 cells, using NUSAP1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using NUSAP1 antibody.

**Product Name: ANKT Rabbit Polyclonal Antibody**  
**Catalog #: APRab06907**

---



Western Blot analysis of various cells using ANKT Polyclonal Antibody

### **Note**

For research use only.