

Product Name: ATP5H Rabbit Polyclonal Antibody
Catalog #: APRab07334



Summary

Production Name	ATP5H Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	ATP5H
Alternative Names	ATP5H; My032; ATP synthase subunit d; mitochondrial; ATPase subunit d
Gene ID	10476.0
SwissProt ID	O75947.The antiserum was produced against synthesized peptide derived from human ATP5H. AA range:111-160

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:20000..
Molecular Weight	26kD

Background

Product Name: ATP5H Rabbit Polyclonal Antibody
Catalog #: AP Rab07334




Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F₁, and the membrane-spanning component, F_o, which comprises the proton channel. The F₁ complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The F_o seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the d subunit of the F_o complex. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. In addition, three pseudogenes are located on chromosomes 9, 12 and 15. [provided by RefSeq, Jun 2010],function:Mitochondrial membrane ATP synthase (F₁)F₀ ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core, and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F(0) domain and the peripheric stalk, which acts as a stator to hold the catalytic alpha(3)beta(3) subcomplex and subunit a/ATP6 static relative to the rotary elements.,similarity:Belongs to the ATPase d subunit family.,subunit:F-type ATPases have 2 components, CF(1) - the catalytic core - and CF(0) - the membrane proton channel. CF(0) seems to have nine subunits: a, b, c, d, e, f, g, F6 and 8 (or A6L),

Research Area


Oxidative phosphorylation;Alzheimer's disease;Parkinson's disease;Huntington's disease;

Image Data



Western blot analysis of lysates from HepG2 and Jurkat cells, using ATP5H Antibody. The lane on the right is blocked with the synthesized peptide.

Product Name: ATP5H Rabbit Polyclonal Antibody
Catalog #: APRab07334



Western blot analysis of the lysates from HepG2 cells using ATP5H antibody.



Western Blot analysis of various cells using ATP5H Polyclonal Antibody diluted at 1: 2000

Note

For research use only.