

Summary

Production Name	ATP5J2 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	ATP5J2	
Alternative Names	ATP synthase f chain mitochondrial; ATP5JL; ATPK	
Gene ID	9551.0	
SwissProt ID	A6ND55.The antiserum was produced against synthesized peptide derived from	
	human ATP5J2. AA range:21-70	

Application

Dilution Ratio	IHC 1:100-1:300	ELISA: 1:5000
Dilution Ratio	IHC 1:100-1:300	ELISA: 1:5000

Molecular Weight

Background

Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner

Product Name: ATP5J2 Rabbit Polyclonal Antibody Catalog #: APRab07337



membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The catalytic portion of mitochondrial ATP synthase consists of five different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and single representatives of the gamma, delta, and epsilon subunits. The proton channel likely has nine subunits (a, b, c, d, e, f, g, F6 and 8). ATP5J2 (ATP synthase, H+ transporting, mitochondrial Fo complex subunit F2) encodes the f subunit of the Fo complex. Alternatively spliced transcript variants encoding different isoforms have been identified for ATP5J2. ATP5J2 has multiple pseudogenes. Naturally occurring read-through transcription also exists between ATP5J2 and the downstream pentatricopeptide repeat domain 1 (PTCD1) gene. purine nucleotide metabolic process, purine nucleotide biosynthetic process, ATP biosynthetic process, ion transport, cation transport, hydrogen transport, nucleoside triphosphate metabolic process, nucleoside triphosphate biosynthetic process, purine nucleoside triphosphate metabolic process, purine nucleoside triphosphate biosynthetic process, purine ribonucleotide metabolic process, purine ribonucleotide biosynthetic process, nucleotide biosynthetic process, ribonucleoside triphosphate metabolic process, ribonucleoside triphosphate biosynthetic process, purine ribonucleoside triphosphate metabolic process, purine ribonucleoside triphosphate biosynthetic process, ribonucleotide metabolic process, ribonucleotide biosynthetic process, monovalent inorganic cation transport, proton transport, nucleobase, nucleoside and nucleotide biosynthetic process, nucleobase, nucleoside, nucleotide and nucleic acid biosynthetic process, nitrogen compound biosynthetic process, ATP metabolic process,

Research Area

Oxidative phosphorylation;

Image Data



Immunohistochemistry analysis of ATP5J2 antibody in paraffin-embedded human lung carcinoma tissue.

Note

For research use only.