

Product Name: ATP-citrate synthase Rabbit Polyclonal Antibody
Catalog #: APRab07349



Summary

Production Name	ATP-citrate synthase Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat,Monkey

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	ACLY
Alternative Names	ACLY; ATP-citrate synthase; ATP-citrate; pro-S-)-lyase; ACL; Citrate cleavage enzyme
Gene ID	47.0
SwissProt ID	P53396.The antiserum was produced against synthesized peptide derived from human ATP-Citrate Lyase. AA range:420-469

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Molecular Weight	120kD

Product Name: ATP-citrate synthase Rabbit Polyclonal Antibody
Catalog #: AP Rab07349




Background

ATP citrate lyase (ACLY) Homo sapiens ATP citrate lyase is the primary enzyme responsible for the synthesis of cytosolic acetyl-CoA in many tissues. The enzyme is a tetramer (relative molecular weight approximately 440,000) of apparently identical subunits. It catalyzes the formation of acetyl-CoA and oxaloacetate from citrate and CoA with a concomitant hydrolysis of ATP to ADP and phosphate. The product, acetyl-CoA, serves several important biosynthetic pathways, including lipogenesis and cholesterologenesis. In nervous tissue, ATP citrate-lyase may be involved in the biosynthesis of acetylcholine. Multiple transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Dec 2014], catalytic activity: ADP + phosphate + acetyl-CoA + oxaloacetate = ATP + citrate + CoA, function: ATP citrate-lyase is the primary enzyme responsible for the synthesis of cytosolic acetyl-CoA in many tissues. Has a central role in de novo lipid synthesis. In nervous tissue it may be involved in the biosynthesis of acetylcholine, similarity: In the C-terminal section; belongs to the succinate/malate CoA ligase alpha subunit family, similarity: In the N-terminal section; belongs to the succinate/malate CoA ligase beta subunit family, subunit: Homotetramer,

Research Area


Citrate cycle (TCA cycle);

Image Data




Western blot analysis of lysates from COS7 cells, treated with Calyculin 50nM 30', using ATP-Citrate Lyase Antibody. The lane on the right is blocked with the synthesized peptide.

Product Name: ATP-citrate synthase Rabbit Polyclonal Antibody
Catalog #: AP Rab07349



Western Blot analysis of various cells using ATP-citrate synthase Polyclonal Antibody diluted at 1 : 1000



Western Blot analysis of COS7 cells using ATP-citrate synthase Polyclonal Antibody diluted at 1 : 1000

Note

For research use only.