

Summary

Production Name	BRSK1 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	BRSK1
Alternative Names	BRSK1; KIAA1811; SAD1; SADB; Serine/threonine-protein kinase BRSK1; Brain-selective
	kinase 1; Brain-specific serine/threonine-protein kinase 1; BR serine/threonine-protein
	kinase 1; Serine/threonine-protein kinase SAD-B; Synapses of Amphids
Gene ID	84446.0
SwissProt ID	Q8TDC3.The antiserum was produced against synthesized peptide derived from human
	BRSK1. AA range:361-410

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications.
Molecular Weight	87kD



Background

catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Thr-205 by STK11 in complex with STE20-related adapter-alpha (STRAD alpha) pseudo kinase and CAB39., function: Required for the polarization of forebrain neurons which endows axons and dendrites with distinct properties, possibly by locally regulating phosphorylation of microtubule-associated proteins (By similarity). May be involved in the regulation of G2/M arrest in response to UV- or methyl methane sulfonate (MMS)-induced, but not IRinduced, DNA damage. Phosphorylates WEE1 and CDC25B in vitro and CDC25C in vitro and in vivo., similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. AMPK subfamily..similarity:Contains 1 protein kinase domain.,similarity:Contains 1 UBA domain.,subcellular location:Nuclear in the absence of DNA damage. Translocated to the nucleus in response to UV- or MMS-induced DNA damage., tissue specificity: Widely expressed, with highest levels in brain and testis. Protein levels remain constant throughout the cell cycle., catalytic activity: ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Activated by phosphorylation on Thr-205 by STK11 in complex with STE20-related adapter-alpha (STRAD alpha) pseudo kinase and CAB39, function: Required for the polarization of forebrain neurons which endows axons and dendrites with distinct properties, possibly by locally regulating phosphorylation of microtubule-associated proteins (By similarity). May be involved in the regulation of G2/M arrest in response to UV- or methyl methane sulfonate (MMS)-induced, but not IR-induced, DNA damage. Phosphorylates WEE1 and CDC25B in vitro and CDC25C in vitro and in vivo., similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. AMPK subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 1 UBA domain.,subcellular location:Nuclear in the absence of DNA damage. Translocated to the nucleus in response to UV- or MMS-induced DNA damage, tissue specificity: Widely expressed, with highest levels in brain and testis. Protein levels remain constant throughout the cell cycle.,

Research Area

Image Data



Product Name: BRSK1 Rabbit Polyclonal Antibody Catalog #: APRab07668



Western blot analysis of lysates from 293 cells, using BRSK1 Antibody. The lane on the right is blocked with the synthesized



Western blot analysis of the lysates from COLO205 cells using BRSK1 antibody.



Western Blot analysis of various cells using BRSK1 Polyclonal Antibody

Note

For research use only.