# Product Name: DHA Kinase Rabbit Polyclonal Antibody Catalog #: APRab09958

## **Summary**

Production Name DHA Kinase Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

**Application** WB,IHC,ELISA **Reactivity** Human,Mouse,Rat

## **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

#### **Immunogen**

Storage

Gene Name DAK

Alternative Names DAK; Bifunctional ATP-dependent dihydroxyacetone kinase/FAD-AMP lyase; cyclizing

**Gene ID** 26007.0

Q3LXA3. The antiserum was produced against synthesized peptide derived from human

DAK. AA range:91-140

# **Application**

**SwissProt ID** 

**Dilution Ratio** WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000...

Molecular Weight 59kD

# **Background**




This gene is a member of the family of dihydroxyacetone kinases, which have a protein structure distinct from other kinases. The product of this gene phosphorylates dihydroxyacetone, and also catalyzes the formation of riboflavin 4',5'-phosphate (aka cyclin FMN) from FAD. Several alternatively spliced transcript variants have been identified, but the full-length nature of only one has been determined. [provided by RefSeq, Jul 2008],catalytic activity:ATP + glycerone = ADP + glycerone phosphate.,catalytic activity:FAD = AMP + riboflavin cyclic-4',5'phosphate,,cofactor:Magnesium,,cofactor:Manganese or cobalt; for FAD-AMP lyase activity,,enzyme regulation:Each activity is inhibited by the substrate(s) of the other, function: Catalyzes both the phosphorylation of dihydroxyacetone and the splitting of ribonucleoside diphosphate-X compounds among which FAD is the best substrate, similarity: Belongs to the dihydroxyacetone kinase (DAK) family, similarity: Contains 1 DAK1 (dihydroxyacetone kinase subunit 1) domain., similarity: Contains 1 DAK2 (dihydroxyacetone kinase subunit 2) domain., subunit: Homodimer.,


### **Research Area**

Glycerolipid metabolism;RIG-I-like receptor;

## **Image Data**



Western blot analysis of lysates from K562 and A549 cells, using DAK Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded Human prostate cancer. Antibody was diluted at 1:100 (4°,overnight) . High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838 (right) obtaned from antibody was pre-absorbed by immunogen peptide.

#### Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838