

Product Name: ECA39 Rabbit Polyclonal Antibody
Catalog #: APRab10277



Summary

Production Name	ECA39 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	BCAT1
Alternative Names	BCAT1; BCT1; ECA39; Branched-chain-amino-acid aminotransferase, cytosolic; BCAT(c); Protein ECA39
Gene ID	586.0
SwissProt ID	P54687.The antiserum was produced against synthesized peptide derived from the Internal region of human BCAT1. AA range:231-280

Application

Dilution Ratio	WB 1:500 - 1:2000. IHC-p: 1:100-1:300. ELISA: 1:20000..
Molecular Weight	43kD

Background

branched chain amino acid transaminase 1 (BCAT1) Homo sapiens This gene encodes the cytosolic form of the enzyme branched-chain amino acid transaminase. This enzyme catalyzes the reversible transamination of branched-chain alpha-keto acids to branched-chain L-amino acids essential for cell growth. Two different clinical disorders have been attributed to a defect of branched-chain amino acid transamination: hypervalinemia and hyperleucine-isoleucinemia. As there is also a gene encoding a mitochondrial form of this enzyme, mutations in either gene may contribute to these disorders. Alternatively spliced transcript variants have been described. [provided by RefSeq, May 2010], catalytic activity: 2-oxoglutaric acid + L-isoleucine = (S)-3-methyl-2-oxopentanoic acid + L-glutamic acid, catalytic activity: 2-oxoglutaric acid + L-valine = 3-methyl-2-oxobutanoic acid + L-glutamic acid, catalytic activity: L-leucine + 2-oxoglutarate = 4-methyl-2-oxopentanoate + L-glutamate, cofactor: Pyridoxal phosphate, function: Catalyzes the first reaction in the catabolism of the essential branched chain amino acids leucine, isoleucine, and valine, similarity: Belongs to the class-IV pyridoxal-phosphate-dependent aminotransferase family, subunit: Homodimer, tissue specificity: During embryogenesis, expressed in the brain and kidney. Overexpressed in C-myc induced tumors such as Burkitt's lymphoma,


Research Area

Valine; leucine and isoleucine degradation; Valine; leucine and isoleucine biosynthesis; Pantothenate and CoA biosynthesis;

Image Data



Western Blot analysis of K562 cells using ECA39 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Product Name: ECA39 Rabbit Polyclonal Antibody
Catalog #: APRab10277



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100

Note

For research use only.