

Summary

Production Name	EP-CAM Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human, Rat, Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	EPCAM
Alternative Names	EPCAM; GA733-2; M1S2; M4S1; MIC18; TACSTD1; TROP1; Epithelial cell adhesion
	molecule; Ep-CAM; Adenocarcinoma-associated antigen; Cell surface glycoprotein
	Trop-1; Epithelial cell surface antigen; Epithelial glycoprotein; EGP; Epithelial gly
Gene ID	4072.0
SwissProt ID	P16422.The antiserum was produced against synthesized peptide derived from human
	TACD1. AA range:116-165

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications.
Molecular Weight	34kD




Background


This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy. [provided by RefSeq, Dec 2008],

Research Area

Image Data




Western blot analysis of lysates from Jurkat, HUVEC, and COLO205 cells, using TACD1 Antibody. The lane on the right is



Western Blot analysis of various cells using EP-CAM Polyclonal Antibody diluted at 1: 2000

Product Name: EP-CAM Rabbit Polyclonal Antibody Catalog #: APRab10508





Western Blot analysis of 3T3 cells using EP-CAM Polyclonal Antibody diluted at 1: 2000

Note

For research use only.