

**Product Name: FAT10 Rabbit Polyclonal Antibody**  
**Catalog #: APRab10846**



## Summary

|                        |                                  |
|------------------------|----------------------------------|
| <b>Production Name</b> | FAT10 Rabbit Polyclonal Antibody |
| <b>Description</b>     | Rabbit Polyclonal Antibody       |
| <b>Host</b>            | Rabbit                           |
| <b>Application</b>     | WB,ELISA                         |
| <b>Reactivity</b>      | Human,Rat,Mouse                  |

## Performance

|                     |  |
|---------------------|--|
| <b>Conjugation</b>  | Unconjugated   |
| <b>Modification</b> | Unmodified   |
| <b>Isotype</b>      | IgG  |
| <b>Clonality</b>    | Polyclonal   |
| <b>Form</b>         | Liquid   |
| <b>Storage</b>      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |
| <b>Buffer</b>       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.       |
| <b>Purification</b> | Affinity purification  |

## Immunogen

|                          |   |
|--------------------------|---|
| <b>Gene Name</b>         | UBD   |
| <b>Alternative Names</b> | UBD; FAT10; Ubiquitin D; Diubiquitin; Ubiquitin-like protein FAT10          |
| <b>Gene ID</b>           | 10537.0   |
| <b>SwissProt ID</b>      | O15205.Synthesized peptide derived from the Internal region of human FAT10. |

## Application

|                         |                                   |
|-------------------------|-----------------------------------|
| <b>Dilution Ratio</b>   | WB 1:500 - 1:2000. ELISA: 1:10000 |
| <b>Molecular Weight</b> | 18kD                              |

## Background

similarity:Contains 2 ubiquitin-like domains.,subunit:Interacts with MAD2.,similarity:Contains 2 ubiquitin-like


**Product Name: FAT10 Rabbit Polyclonal Antibody**  
**Catalog #: APRab10846**




domains.,subunit:Interacts with MAD2.,

## Research Area

## Image Data



Western blot analysis of K562 293T using FAT10 antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000



Western blot analysis of K562 using FAT10 antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000

## Note

For research use only.