

Summary

| Production Name | FGF-22 Rabbit Polyclonal Antibody |
|-----------------|-----------------------------------|
| Description | Rabbit Polyclonal Antibody |
| Host | Rabbit |
| Application | IF,IHC,WB, |
| Reactivity | Human, Mouse, Rat |

Performance

| Conjugation | Unconjugated |
|--------------|--|
| Modification | Unmodified |
| lsotype | lgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw |
| | cycles. |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification |

Immunogen

| Gene Name | FGF22 |
|-------------------|--|
| Alternative Names | FGF22; Fibroblast growth factor 22; FGF-22 |
| Gene ID | 27006.0 |
| SwissProt ID | Q9HCT0.The antiserum was produced against synthesized peptide derived from |
| | human FGF22. AA range:71-120 |

Application

| Dilution Ratio | WB 1:500 - 1:2000 IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:40000. Not yet tested |
|------------------|--|
| | in other applications. |
| Molecular Weight | 20kD |



Background

The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities and are involved in a variety of biological processes including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. The mouse homolog of this gene was found to be preferentially expressed in the inner root sheath of the hair follicle, which suggested a role in hair development. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2014],function:May be involved in hair development.,similarity:Belongs to the heparin-binding growth factors family.,subunit:Interacts with FGFBP1.,


Research Area

MAPK_ERK_Growth;MAPK_G_Protein;Regulates Actin and Cytoskeleton;Pathways in cancer;Melanoma;

Image Data




Immunofluorescence analysis of MCF7 cells, using FGF22 Antibody. The picture on the right is blocked with the synthesized




Immunohistochemistry analysis of paraffin-embedded human ovary tissue, using FGF22 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from Jurkat and HT-29 cells, using FGF22 Antibody. The lane on the right is blocked with the



Western Blot analysis of various cells using FGF-22 Polyclonal Antibody diluted at 1: 1000

Note

For research use only.