Product Name: Gremlin-2 Rabbit Polyclonal Antibody Catalog #: APRab11757



Summary

Production Name Gremlin-2 Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IHC,ELISA **Reactivity** Human,Mouse

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

ClonalityPolyclonalFormLiquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name GREM2 CKTSF1B2 DAND3 PRDC

Gremlin-2 (Cysteine knot superfamily 1, BMP antagonist 2;DAN domain family member Alternative Names

3;Protein related to DAN and cerberus)

Gene ID 64388.0

Q9H772.The antiserum was produced against synthesized peptide derived from the **SwissProt ID**

Internal region of human GREM2. AA range:71-120

Application

Dilution Ratio WB 1:500-2000,IHC 1:500-200, ELISA 1:10000-20000.

Molecular Weight 19kD

Product Name: Gremlin-2 Rabbit Polyclonal Antibody Catalog #: APRab11757




Background

This gene encodes a member of the BMP (bone morphogenic protein) antagonist family. Like BMPs, BMP antagonists contain cystine knots and typically form homo- and heterodimers. The CAN (cerberus and dan) subfamily of BMP antagonists, to which this gene belongs, is characterized by a C-terminal cystine knot with an eight-membered ring. The antagonistic effect of the secreted glycosylated protein encoded by this gene is likely due to its direct binding to BMP proteins. As an antagonist of BMP, this gene may play a role in regulating organogenesis, body patterning, and tissue differentiation. [provided by RefSeq, Jul 2008],function:Cytokine that inhibits the activity of BMP2 and BMP4 in a dose-dependent manner. Antagonized BMP4-induced suppression of progesterone production in granulosa cells.,similarity:Belongs to the DAN family.,similarity:Contains 1 CTCK (C-terminal cystine knot-like) domain.,subunit:Interacts with BMP2 and BMP4.,


Research Area

Image Data



1, mouse-kidney 2, 293T 3, mouse-lung


Western blot analysis of RAT-brain lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human-small-intestine, antibody was diluted at 1:200

Catalog #: APRab11757






Immunohistochemical analysis of paraffin intestine, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-colon-cancer, antibody was diluted at 1:200

Note

For research use only.