

### Summary

Production Name	hnRNP DL Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human, Mouse, Rat

### Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

### Immunogen

Gene Name	HNRPDL	
Alternative Names	HNRPDL; JKTBP; Heterogeneous nuclear ribonucleoprotein D-like; hnRNP D-like;	
	hnRNP DL; AU-rich element RNA-binding factor; JKT41-binding protein; Protein laAUF1	
Gene ID	9987.0	
SwissProt ID	O14979.The antiserum was produced against synthesized peptide derived from human	
	HNRPDL. AA range:241-290	

# Application

Dilution Ratio	IHC 1:100-1:300	ELISA: 1:20000

**Molecular Weight** 

## Background


# Catalog #: APRab12142



heterogeneous nuclear ribonucleoprotein D like(HNRNPDL) Homo sapiens This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two RRM domains that bind to RNAs. Three alternatively spliced transcript variants have been described for this gene. One of the variants is probably not translated because the transcript is a candidate for nonsense-mediated mRNA decay. The protein isoforms encoded by this gene are similar to its family memberfunction: Acts as a transcriptional regulator. Promotes transcription repression. Promotes transcription activation in differentiated myotubes (By similarity). Binds to double- and single-stranded DNA sequences. Binds to the transcription suppressor CATR sequence of the COX5B promoter (By similarity). Binds with high affinity to RNA molecules that contain AU-rich elements (AREs) found within the 3'-UTR of many proto-oncogenes and cytokine mRNAs. Binds both to nuclear and cytoplasmic poly(A) mRNAs. Binds to poly(G) and poly(A), but not to poly(U) or poly(C) RNA homopolymers. Binds to the 5'-ACUAGC-3' RNA consensus sequence., induction: Up-regulated by 12-Otetradecanoylphorbol 13-acetate (TPA) in macrophages and retinoic acid (RA) in granulocytes (at protein level). Downregulated by interleukin-4 (IL-4), PTM:Dimethylation of Arg-408 is probably of the asymmetric type, similarity:Contains 2 RRM (RNA recognition motif) domains, subcellular location: Shuttles between the nucleus and the cytoplasm in a TNPO1dependent manner., subunit: Interacts with ZNF148 (By similarity). Interacts with TNPO1., tissue specificity: Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine, colon and leukocytes. Expressed in myeloid leukemia, gastric adenocarcinoma, cervical carcinoma, hepatoma, fibrosarcoma, colon adenocarcinoma, epidermoid carcinoma, osteosarcoma and urinary bladder carcinoma cells.,

### **Research Area**

### Image Data



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using HNRPDL Antibody. The picture

# Product Name: hnRNP DL Rabbit Polyclonal Antibody Catalog #: APRab12142



on the right is blocked with the synthesized peptide.

#### Note

For research use only.