

Product Name: Insulin Rabbit Polyclonal Antibody
Catalog #: APRab12638



Summary

Production Name	Insulin Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	INS
Alternative Names	INS; Insulin
Gene ID	3630.0
SwissProt ID	P01308.The antiserum was produced against synthesized peptide derived from human Insulin. AA range:49-98

Application

Dilution Ratio	IHC 1:100-1:300 ELISA: 1:10000
Molecular Weight	

Background

After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three peptides: the B chain and

Product Name: Insulin Rabbit Polyclonal Antibody
Catalog #: AP Rab12638




A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2010],disease:Defects in INS are the cause of familial hyperproinsulinemia [MIM:176730],function:Insulin decreases blood glucose concentration. It increases cell permeability to monosaccharides, amino acids and fatty acids. It accelerates glycolysis, the pentose phosphate cycle, and glycogen synthesis in liver.,function:Preptin undergoes glucose-mediated co-secretion with insulin, and acts as physiological amplifier of glucose-mediated insulin secretion. Exhibits osteogenic properties by increasing osteoblast mitogenic activity through phosphoactivation of MAPK1 and MAPK3,function:The insulin-like growth factors possess growth-promoting activity. In vitro, they are potent mitogens for cultured cells. IGF-II is influenced by placental lactogen and may play a role in fetal development.,mass spectrometry: PubMed:12586351; PubMed:15359740,online information:Clinical information on Eli Lilly insulin products,online information:Insulin entry,online information:Insulin-like growth factor 2 entry,online information:Protein of the 20th century - Issue 9 of April 2001,pharmaceutical:Available under the names Humulin or Humalog (Eli Lilly) and Novolin (Novo Nordisk). Used in the treatment of diabetes. Humalog is an insulin analog with 52-Lys-Pro-53 instead of 52-Pro-Lys-53.,similarity:Belongs to the insulin family.,subunit:Heterodimer of a B chain and an A chain linked by two disulfide bonds.,

Research Area

Oocyte meiosis;Regulation of autophagy;mTOR;Regulates Actin and Cytoskeleton;Insulin_Receptor;Progesterone-mediated oocyte maturation;Type II diabetes mellitus;Type I diabetes mellitus;Maturity onset diabetes of the young;Aldosterone-regulated sodium reabsorption;Prostate cancer;

Image Data



Immunohistochemistry analysis of Insulin antibody in paraffin-embedded human pancreas tissue.

Note

For research use only.

Product Name: Insulin Rabbit Polyclonal Antibody
Catalog #: APRab12638

