

# Summary

Production Name	KDEL Receptor 2 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IF,WB,IHC
Reactivity	Human,Mouse,Rat

### Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

### Immunogen

Gene Name	KDELR2
Alternative Names	KDELR2; ERD2.2; ER lumen protein retaining receptor 2; ERD2-like protein 1; ELP-1;
	KDEL endoplasmic reticulum protein retention receptor 2; KDEL receptor 2
Gene ID	11014.0
SwissProt ID	P33947.The antiserum was produced against synthesized peptide derived from human
	ERD22. AA range:81-130

# Application

Dilution Ratio	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested
	in other applications.

## Product Name: KDEL Receptor 2 Rabbit Polyclonal Antibody Catalog #: APRab12960



Molecular Weight 24kD


### Background

KDEL endoplasmic reticulum protein retention receptor 2(KDELR2) Homo sapiens Retention of resident soluble proteins in the lumen of the endoplasmic reticulum (ER) is achieved in both yeast and animal cells by their continual retrieval from the cis-Golgi, or a pre-Golgi compartment. Sorting of these proteins is dependent on a C-terminal tetrapeptide signal, usually lys-asp-glu-leu (KDEL) in animal cells, and his-asp-glu-leu (HDEL) in S. cerevisiae. This process is mediated by a receptor that recognizes, and binds the tetrapeptide-containing protein, and returns it to the ER. In yeast, the sorting receptor encoded by a single gene, ERD2, is a seven-transmembrane protein. Unlike yeast, several human homologs of the ERD2 gene, constituting the KDEL receptor gene family, have been described. KDELR2 was the second member of the family to be identified, and it encodes a protein which is 83% identical to the KDELR1 gene product. Alternative splicing rfunction:Required for the retention of luminal endoplasmic reticulum proteins. Determines the specificity of the luminal ER protein retention system. Also required for normal vesicular traffic through the Golgi. This receptor recognizes K-D-E-L,similarity:Belongs to the ERD2 family.,

#### **Research Area**


Vibrio cholerae infection;

## Image Data




Immunofluorescence analysis of A549 cells, using ERD22 Antibody. The picture on the right is blocked with the synthesized peptide.






Western blot analysis of lysates from HUVEC and COLO cells, using ERD22 Antibody. The lane on the right is blocked with



Western Blot analysis of various cells using KDEL Receptor 2 Polyclonal Antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded Human lung cancer. Antibody was diluted at 1:100 (4°,overnight) . High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

#### Note

For research use only.