

Summary

Production Name	LTB4DH Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	PTGR1
Alternative Names	PTGR1; LTB4DH; Prostaglandin reductase 1; PRG-1; 15-oxoprostaglandin 13-reductase; NADP-dependent leukotriene B4 12-hydroxydehydrogenase
Gene ID	22949.0
SwissProt ID	Q14914.The antiserum was produced against synthesized peptide derived from human PTGR1. AA range:75-124

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications.
Molecular Weight	36kD

Product Name: LTB4DH Rabbit Polyclonal Antibody
Catalog #: AP Rab13472




Background


This gene encodes an enzyme that is involved in the inactivation of the chemotactic factor, leukotriene B4. The encoded protein specifically catalyzes the NADP+ dependent conversion of leukotriene B4 to 12-oxo-leukotriene B4. A pseudogene of this gene is found on chromosome 1. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2009], catalytic activity: 11- α -hydroxy-9,15-dioxoprost-5-enoate + NAD(P)(+) = (5Z)-(13E)-11- α -hydroxy-9,15-dioxoprost-5,13-dienoate + NAD(P)H., catalytic activity: n-alkanal + NAD(P)(+) = alk-2-enal + NAD(P)H., function: Functions as 15-oxo-prostaglandin 13-reductase and acts on 15-oxo-PGE1, 15-oxo-PGE2 and 15-oxo-PGE2- α . Has no activity towards PGE1, PGE2 and PGE2- α (By similarity). Catalyzes the conversion of leukotriene B4 into its biologically less active metabolite, 12-oxo-leukotriene B4. This is an initial and key step of metabolic inactivation of leukotriene B4., similarity: Belongs to the NADP-dependent oxidoreductase L4BD family., subunit: Monomer or homodimer., tissue specificity: High expression in the kidney, liver, and intestine but not in leukocytes.,

Research Area

Image Data




Western blot analysis of lysates from HT-29 cells, using PTGR1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from 293 cells using PTGR1 antibody.

Product Name: LTB4DH Rabbit Polyclonal Antibody
Catalog #: APRab13472



Western Blot analysis of various cells using LTB4DH Polyclonal Antibody

Note

For research use only.