

Summary

Production Name	Manic Fringe Rabbit Polyclonal Antibody	
Description	Rabbit Polyclonal Antibody	
Host	Rabbit	
Application	WB	
Reactivity	Human, Mouse, Rat	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	MFNG
Alternative Names	MFNG; Beta-1; 3-N-acetylglucosaminyltransferase manic fringe; O-fucosylpeptide 3-
	beta-N-acetylglucosaminyltransferase
Gene ID	4242.0
SwissProt ID	O00587.The antiserum was produced against synthesized peptide derived from human
	MFNG. AA range:61-110

Application

Dilution Ratio	WB 1:500-1:2000. ELISA: 1:5000.
Molecular Weight	38kD

Background

Product Name: Manic Fringe Rabbit Polyclonal Antibody Catalog #: APRab13619


This gene is a member of the fringe gene family which also includes radical and lunatic fringe genes. They all encode evolutionarily conserved secreted proteins that act in the Notch receptor pathway to demarcate boundaries during embryonic development. While their genomic structure is distinct from other glycosyltransferases, fringe proteins have a fucose-specific beta-1,3-N-acetylglucosaminyltransferase activity that leads to elongation of O-linked fucose residues on Notch, which alters Notch signaling. [provided by RefSeq, Oct 2009],catalytic activity:Transfers a beta-D-GlcNAc residue from UDP-D-GlcNAc to the fucose residue of a fucosylated protein acceptor.,function:Glycosyltransferase involved in the elongation of O-linked ligands to activate Notch signaling. Possesses fucose-specific beta-1,3-N-acetylglucosaminyltransferase activity.,online information:Beta-1,3-N-acetylglucosaminyltransferase manic fringe,online information:GlycoGene database,similarity:Belongs to the glycosyltransferase 31 family.,

EnkiLife

Research Area

Notch;

Image Data



Western blot analysis of the lysates from HepG2 cells using MFNG antibody.

Note

For research use only.