

**Product Name: mGluR-4 Rabbit Polyclonal Antibody**  
**Catalog #: APRab13861**



## Summary

<b>Production Name</b>	mGluR-4 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	IHC, WB,
<b>Reactivity</b>	Human, Mouse, Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	GRM4
<b>Alternative Names</b>	GRM4; GPRC1D; MGLUR4; Metabotropic glutamate receptor 4; mGluR4
<b>Gene ID</b>	2914.0
<b>SwissProt ID</b>	Q14833. The antiserum was produced against synthesized peptide derived from human mGluR4. AA range: 851-900

## Application

<b>Dilution Ratio</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000..
<b>Molecular Weight</b>	100kD

## Background

**Product Name: mGluR-4 Rabbit Polyclonal Antibody**  
**Catalog #: APRab13861**




glutamate metabotropic receptor 4 (GRM4) Homo sapiens L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Several transcript variants encoding different isoforms have been found for this function: Receptor for glutamate. The activity of this receptor is mediated by a G-protein that inhibits adenylate cyclase activity., similarity: Belongs to the G-protein coupled receptor 3 family., subunit: Interacts with PICK1., tissue specificity: Strongly expressed in the cerebellum. Expressed at low levels in hippocampus, hypothalamus and thalamus. No expression detected in liver.,


## Research Area

Neuroactive ligand-receptor interaction; Taste transduction;

## Image Data




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using mGluR4 Antibody. The picture on the right is blocked with the synthesized peptide.




**Product Name: mGluR-4 Rabbit Polyclonal Antibody**  
**Catalog #: APRab13861**



Western blot analysis of lysates from 293 cells, treated with Forskolin 40nM 30', using mGluR4 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using mGluR-4 Polyclonal Antibody diluted at 1: 500



Western Blot analysis of 293 cells using mGluR-4 Polyclonal Antibody diluted at 1: 500

**Note**

For research use only.