## **Product Name: MIXL1 Rabbit Polyclonal Antibody**

Catalog #: APRab13922



## **Summary**

Production Name MIXL1 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

Host Rabbit
Application WB

Reactivity Human, Mouse

#### **Performance**

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Purification	Affinity purification

#### **Immunogen**

Gene Name MIXL1 MIXL

**Alternative Names** 

**Gene ID** 83881.0

**SwissProt ID** Q9H2W2.Synthesized peptide derived from human protein . at AA range: 90-170

## **Application**

**Dilution Ratio** WB 1:500-2000 ELISA 1:5000-20000

Molecular Weight 25kD

### **Background**

Mix paired-like homeobox(MIXL1) Homo sapiens Homeodomain proteins, such as MIXL1, are transcription factors that regulate cell fate during development (Hart et al., 2005 [PubMed 15982639]).[supplied by OMIM, Mar

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# **Product Name: MIXL1 Rabbit Polyclonal Antibody**


Catalog #: APRab13922



2008], function: Transcription factor that play a central role in proper axial mesendoderm morphogenesis and endoderm formation. Required for efficient differentiation of cells from the primitive streak stage to blood, by acting early in the recruitment and/or expansion of mesodermal progenitors to the hemangioblastic and hematopoietic lineages. Also involved in the morphogenesis of the heart and the gut during embryogenesis. Acts as a negative regulator of brachyury expression., PTM: Phosphorylated at multiple sites., similarity: Belongs to the paired homeobox family., similarity: Contains 1 homeobox DNA-binding domain., tissue specificity: Restricted to progenitors and secondary lymph tissues. In normal hematopoiesis, it is restricted to immature B-and T-lymphoid cells. Present in differentiating embryonic stem cells (at protein level).,

#### **Research Area**

#### **Image Data**



Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4° over night

#### Note

For research use only.