

Summary

Production Name	MTA1 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	MTA1
Alternative Names	MTA1; Metastasis-associated protein MTA1
Gene ID	9112.0
SwissProt ID	Q13330.The antiserum was produced against synthesized peptide derived from human MTA1. AA range:171-220

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:40000
Molecular Weight	80kD

Background


Product Name: MTA1 Rabbit Polyclonal Antibody
Catalog #: APRab14198



This gene encodes a protein that was identified in a screen for genes expressed in metastatic cells, specifically, mammary adenocarcinoma cell lines. Expression of this gene has been correlated with the metastatic potential of at least two types of carcinomas although it is also expressed in many normal tissues. The role it plays in metastasis is unclear. It was initially thought to be the 70kD component of a nucleosome remodeling deacetylase complex, NuRD, but it is more likely that this component is a different but very similar protein. These two proteins are so closely related, though, that they share the same types of domains. These domains include two DNA binding domains, a dimerization domain, and a domain commonly found in proteins that methylate DNA. The profile and activity of this gene product suggest that it is involved in regulating transcription and that this may be accomplished by chromatin remodeling. **Highly expressed in metastatic cells.** **function:** May be involved in the regulation of gene expression by covalent modification of histone proteins. The long isoform is a corepressor of estrogen receptor (ER). The short isoform binds to ER and sequesters it in the cytoplasm and enhances non-genomic responses of ER. **miscellaneous:** The short isoform contains a Leu-Arg-Ile-Leu-Leu motif (ER binding motif). **similarity:** Contains 1 BAH domain. **similarity:** Contains 1 ELM2 domain. **similarity:** Contains 1 GATA-type zinc finger. **similarity:** Contains 1 SANT domain. **subunit:** Component of the nucleosome-remodeling and histone-deacetylase multiprotein complex (NuRD). Interacts with HDAC1 and ITGB3BP/CENPR. **tissue specificity:** Widely expressed. High expression in brain, ovaries, adrenal glands and virgin mammary glands. Higher in tumors than in adjacent normal tissue from the same individual.


Research Area

Image Data



Western blot analysis of lysates from Jurkat cells, using MTA1 Antibody. The lane on the right is blocked with the synthesized peptide.

Product Name: MTA1 Rabbit Polyclonal Antibody
Catalog #: APRab14198



Western blot analysis of the lysates from K562 cells using MTA1 antibody.

Note

For research use only.